arXiv:1603.06179v1 [math.NT] 20 Mar 2016

ON THE INHOMOGENEOUS SPECTRUM OF PERIOD TWO QUADRATICS

Christopher G. Pinner

Abstract. For an α with period two negative continued fraction expansion:

$$\alpha = [0; a_1, a_2, a_3, ..., a_N, \overline{a, b}]^- = \cfrac{1}{a_1 - \cfrac{1}{a_2 - \cfrac{1}{a_3 - \cdots}}}, \quad 2 \leq a < b,$$

we give a complete description of the values in its inhomogeneous Lagrange spectrum

$$\mathbf{L}(\alpha) := \{M(\alpha, \gamma) \ : \ \gamma \notin \mathbf{Z} + \alpha\mathbf{Z}\}, \quad M(\alpha, \gamma) := \liminf_{n \to \infty} |n|||n\alpha - \gamma||$$

down to the first limit point. The largest point in the spectrum is always isolated (as conjectured by Barnes and Swinnerton-Dyer).

If a is odd then the first limit point is usually the second or third largest value (the fourth in two cases $(a, b) = (5, 7)$ or $(7, 9)$, and a limit from above when $(a, b) = (3, 4), (3, 5), (3, 6)$). When a is even and b is odd there are infinitely many values above the first limit point. This is also the case when a and b are both even except for $b = 2a - 2$, $b = a + 2$ or $b = a + 4$, $a \geq 4$, when the limit point is usually the third largest value.

1. Introduction.

For a pair of real numbers α and γ, with α irrational, one defines the inhomogeneous approximation constant

$$M(\alpha, \gamma) := \liminf_{n \to \infty} |n|||n\alpha - \gamma||.$$

Of course $M(\alpha, \gamma) = M(\alpha, n\alpha + m \pm \gamma)$ for any n, m in $\mathbf{Z}$. Fixing $\gamma = 0$ and varying α gives the (reciprocal of the) traditional Lagrange spectrum

$$\mathbf{L}^{-1} = \{M(\alpha, 0) \ : \ \alpha \in \mathbf{R}\}.$$

Typeset by $\mathcal{AMS}$-TEX

Here we instead fix α and vary γ to obtain the inhomogeneous Lagrange spectrum for α

$$\mathbf{L}(\alpha) := \{M(\alpha,\gamma) \ : \ \gamma \notin \mathbf{Z}+\alpha\mathbf{Z}\}.$$

For the classical Lagrange spectrum old results of Markoff and Lagrange (see [6]) give a complete description of $\mathbf{L}^{-1}$ from the largest value $1/\sqrt{5}$ down to the first limit point $1/3$. We suppose here that

$$\alpha = [0; a_1, ..., a_{N-1}, \overline{a,b}]^- = \cfrac{1}{a_1 - \cfrac{1}{a_2 - \cfrac{1}{a_3 - \cdots}}}, \quad 2 \le a < b, \tag{1.1}$$

has a period two negative expansion and give a similarly complete description of $\mathbf{L}(\alpha)$ above the first limit point. We dealt with the period one cases $a = b$ in [6]. The first limit point is always a limit from below but need not be a limit point from above, indeed it can even be the second largest point, as was observed for the related inhomogeneous Markoff spectrum [5]. Godwin's example $x^2 - 23y^2$ of a form with non-isolated second minimum corresponds to the special case $a = 5$, $b = 10$ ($r = 10,\ m = 0$ in Theorem 2). It has been conjectured by Barnes–Swinnerton-Dyer [1] that the largest value in $\mathbf{L}(\alpha)$, denoted $\rho(\alpha)$, is isolated for any quadratic α. We show that this is certainly true for all period two α. Many special cases of $\rho(\alpha)$ occur in [1], including the the period two examples $(a,b) = (3,3n)$ and $(a,b) = (2n+1, 2(2n+1))$. We recall (see [1]) that $\mathbf{Z}[\alpha]$ is norm Euclidean with respect to $(x-\alpha y)(x-\bar{\alpha}y) = x^2+bxy+cy^2$ iff $\rho(\alpha) < 1/\sqrt{b^2-4c}$ (see also [2,4]). It is interesting to note that $\sqrt{14} = [4; \overline{4,8}]^-$ (where $\mathbf{Q}(\sqrt{14})$ is a field recently shown by Malcolm Harper to be be Euclidean but not norm-Euclidean (see also [3])) has just one point in its spectrum above $\frac{1}{2\sqrt{14}}$.

Setting

$$\eta := [0; \overline{a,b}]^- = \frac{1}{2}(b - \sqrt{b^2 - 4b/a}),$$
$$\beta := [0; \overline{b,a}]^- = \frac{1}{2}(a - \sqrt{a^2 - 4a/b}),$$

and

$$D := \eta\beta,$$

we define

$$M^*(\alpha,\gamma) := 4(1-D)M(\alpha,\gamma).$$

We write $\rho^*(\alpha) = \rho_1^*(\alpha), \rho_2^*(\alpha), \ldots$ for the successive values of $M^*(\alpha,\gamma)$ above the first accumulation point. For a given α we showed in [6] how to use an α-expansion of γ to evaluate $M(\alpha,\gamma)$. Since $\mathbf{L}(\alpha) = \mathbf{L}(\eta) = \mathbf{L}(\beta)$ it will be convenient to assume throughout that α is purely periodic with $a_{2i-1} = a$ and $a_{2i} = b$ for $i \geq 1$, so that the α-expansions take the form

$$\gamma = \sum_{i=1}^{\infty} (b_{2i-1}\eta + b_{2i}D)D^{i-1}.$$

Defining integers t_i by $b_i = \frac{1}{2}(a_i - 2 + t_i)$ and setting

$$d_i^+ := \sum_{j=0}^{\infty} \left(t_{i+2j+1}\alpha_i + t_{i+2j+2}D\right) D^j,$$
$$d_i^- := \sum_{j=0}^{\infty} \left(t_{i-2j}\alpha_{i-1} + t_{i-2j-1}D\right) D^j,$$

where

$$\alpha_j := \begin{cases} \beta & \text{if } j \text{ is odd,} \\ \eta & \text{if } j \text{ is even,} \end{cases}$$

we showed that if $t_k = a_k$ at most finitely often then

$$M^*(\alpha,\gamma) = \liminf_{k\to\infty} \min\{s_1^*(k), s_2^*(k), s_3^*(k), s_4^*(k)\}$$

with

$$s_1^*(i) = (1 - \alpha_i + d_i^+)(1 - \alpha_{i-1} + d_i^-),$$
$$s_2^*(i) = (1 + \alpha_i - d_i^+)(1 + \alpha_{i-1} + d_i^-),$$
$$s_3^*(i) = (1 - \alpha_i - d_i^+)(1 - \alpha_{i-1} - d_i^-),$$
$$s_4^*(i) = (1 + \alpha_i + d_i^+)(1 + \alpha_{i-1} - d_i^-).$$

When $t_k = a_k$ infinitely often then we need to check the minimum of $s_1^*(k)$ and $s_2^*(k)$ for both both γ and $1-\gamma$. When $t_k \neq a_k$ the expansion of $1 - \alpha - \gamma$ can be obtained

from that of γ by simply replacing the t_i with $-t_i$. Changing a finite number of t_i of course does not affect $M(\alpha,\gamma)$. We also showed in [6] that if $t_k = a_k$ infinitely often with $k \equiv j \bmod 2$ then

$$M^*(\alpha,\gamma) \le \alpha_{j-1} \tag{1.2}$$

(so we only need worry about these when a_j is small) and if $t_i = a_i$ at most finitely often with $|t_k| \ge t$, $k \equiv j \bmod 2$ infinitely often then

$$M^*(\alpha,\gamma) \le (a_j - t)\alpha_{j-1} = 1 - t\alpha_{j-1} + D. \tag{1.3}$$

In view of this last result we would expect the t_i corresponding to the larger values of $M(\alpha,\gamma)$ to be all small (although when the smallest partial quotient a is odd we in fact typically compensate for the impossibilty of $t_{2i-1} = 0$ by selecting the t_{2i} to be roughly b/a away from zero).

We shall define sets S_k of γ with eventually periodic α-expansions b_i. In particular the γ of interest will have their expansions built from the blocks (b_{2k-1}, b_{2k}) of the form

$$\begin{aligned}
A_t &= \left(\frac{1}{2}(a-2), \frac{1}{2}(b-2+t)\right), & A'_t &= \left(\frac{1}{2}(a-2), \frac{1}{2}(b-2-t)\right),\\
B_t &= \left(\frac{1}{2}(a-2-1), \frac{1}{2}(b-2+t)\right), & B'_t &= \left(\frac{1}{2}(a-2+1), \frac{1}{2}(b-2-t)\right),\\
C_t &= \left(\frac{1}{2}(a-2-2), \frac{1}{2}(b-2+t)\right), & C'_t &= \left(\frac{1}{2}(a-2+2), \frac{1}{2}(b-2-t)\right),\\
E_t &= \left(\frac{1}{2}(a-2-3), \frac{1}{2}(b-2+t)\right), & E'_t &= \left(\frac{1}{2}(a-2+3), \frac{1}{2}(b-2-t)\right),\\
F_t &= \left((a-1), \frac{1}{2}(b-2-t)\right), & F'_t &= \left((a-1), \frac{1}{2}(b-2+(t-4))\right).
\end{aligned}$$

2. A complete description down to the first limit point.

The spectrum when $a \ge 4$ is even.

Suppose first that $a \ge 4$ is even.

When b is odd we define the classes of γ

$$S_{-1} = \{\gamma \ : \ \gamma \text{ has periodic expansion, period } A_1A_1A'_1A'_1\},$$

$$S_{-2} = \{\gamma \ : \ \gamma \text{ has periodic expansion, period } C_3\},$$

and for integers $k \geq 0$

$$S_k = \{\gamma \ : \ \gamma \text{ has periodic expansion, period } A_1'A_1(A_1'A_1'A_1A_1)^k\}.$$

For b even we set

$$\begin{aligned}
S_{k,1} &= \{\gamma : \ \gamma \text{ has periodic expansion, period } A_0(A_2{A'}_2)^k\}, \ \ k \geq 0,\\
S_{k,2} &= \{\gamma : \ \gamma \text{ has periodic expansion, period } A_2C_4^kC_2{A'}_2{C'}_4^k{C'}_2\}, \ \ k \geq 1,\\
S_{k,3} &= \{\gamma : \ \gamma \text{ has periodic expansion, period } C_4(C_2C_4)^k\}, \ \ k \geq 1,\\
S_{k,4} &= \{\gamma : \ \gamma \text{ has periodic expansion, period } C_4C_2^k\}, \ \ k \geq 0,\\
S_{k,5} &= \{\gamma : \ \gamma \text{ has periodic expansion, period } A_2C_2^k{A'}_2{C'}_2^k\}, \ \ k \geq 0,\\
S_{-1} &= \{\gamma : \ \gamma \text{ has periodic expansion, period } C_2\},
\end{aligned}$$

and for $(a, b) = (4, 6)$

$$S_{-2} = \{\gamma : \ \gamma \text{ has periodic expansion, period } A_2C_2\},$$

for $(a, b) = (8, 12)$

$$S_{k,6} = \{\gamma : \ \gamma \text{ has periodic expansion, period } (A_2'C_2'A_2C_2)^kA_2'C_2'C_2'A_2C_2C_2\}, \ \ k \geq 0,$$

and for $(a, b) = (6, 10)$

$$S_{k,7} = \{\gamma : \ \gamma \text{ has periodic expansion, period } (C_4C_2)^kA_2'A_2\}, \ \ k \geq 1,$$

We write

$$\delta_\square = M^*(\alpha, \gamma), \ \ \gamma \in S_\square.$$

Theorem 1. *Suppose that $\alpha = [0; \overline{a, b}]$ with $a < b$ and $a \geq 4$ even.*

If b is odd then

$$\rho^*(\alpha) = \begin{cases} \delta_0, & \text{if } b = a + 1,\ a + 3 \text{ or } b \geq 2a - 3,\\ \delta_{-2}, & \text{if } a + 5 \leq b \leq 2a - 5, \end{cases}$$

and the only values of $M^*(\alpha,\gamma)$ *greater than*

$$\delta_\infty := \left(1-\beta-\frac{\beta(1-D)(1+2D^2)}{1+D^2}\right)\left(1-\eta-\frac{D(1-D)}{1+D^2}\right)$$

are δ_{-1}, δ_{-2} *(when* $a+3\le b\le 2a-3$*) and the* $\delta_k \searrow \delta_\infty$, $k\ge 0$.

When b *is even*

$$\rho^*(\alpha)=\delta_{0,1}=(1-\eta)(1-\beta).$$

For $b\ge 2a$ *or* $(a,b)=(4,6)$ *the values of* $M^*(\alpha,\gamma)$ *greater than*

$$\delta_{\infty,1}=\left(1-\eta+\frac{2D^2}{1+D}\right)\left(1-\beta-\frac{2\beta}{1+D}\right)$$

are the $\delta_{k,1}\searrow\delta_{\infty,1}$, $k\ge 0$ *and* $\delta_{0,5}$ *plus* $\delta_{0,4}$ *when* $2a\le b\le 3a-6$, *and* δ_{-2} *when* $(a,b)=(4,6)$.

When $b=2a-2$, $a\ge 8$ *the values of* $M^*(\alpha,\gamma)$ *greater than*

$$\delta_{\infty,2}=\left(1-3\eta+\frac{2D(2-\eta)}{1-D}\right)\left(1+\beta-\frac{2\beta D(1-\eta+D)}{1-D}\right)$$

are $\delta_{0,1}$ *and* $\delta_{0,4}$ *(with the* $\delta_{k,2}\nearrow\delta_{\infty,2}$*).*

When $b=2a-4$, $a\ge 10$ *the values of* $M^*(\alpha,\gamma)$ *greater than*

$$\delta_{\infty,3}=\left(1-\eta-\frac{2\eta D}{1-D}+\frac{2D(1+2D)}{1-D^2}\right)\left(1-3\beta+\frac{2D}{1-D}-\frac{2\beta D(2+D)}{1-D^2}\right)$$

are $\delta_{0,1}$, $\delta_{0,4}$, $\delta_{1,4}$, $\delta_{1,5}$ *and the* $\delta_{k,3}\searrow\delta_{\infty,3}$, $k\ge 1$.

When $b\le 2a-6$ *or* $a=6, b=8$, *the first limit point is*

$$\delta_{4,\infty}=\left(1-\eta+\frac{2D(1-\eta)}{1-D}\right)\left(1-3\beta+\frac{2D(1-\beta)}{1-D}\right)$$

with $\delta_{k,5}\nearrow\delta_{\infty,4}$. *The values above* $\delta_{\infty,4}$ *are* $\delta_{0,1}$, δ_{-1}, *and when* $a+6\le b\le 2a-6$ *the* $\delta_{k,4}\searrow\delta_{\infty,4}$.

For $(a,b)=(8,12)$ *the values above* $\delta_{\infty,6}=\frac{112387809}{2209}\sqrt{138}-\frac{1320256308}{2209}$ *are* $\delta_{0,1}$, $\delta_{1,5}$ *and the* $\delta_{k,6}\searrow\delta_{\infty,6}$, $k\ge 0$.

For $(a,b)=(6,10)$ *the values above* $\delta_{\infty,7}=\frac{6329319}{40}-\frac{6551443}{600}\sqrt{210}$ *are* $\delta_{0,1}$, $\delta_{0,5}$, $\delta_{1,4}$ *with the* $\delta_{k,7}\nearrow\delta_{\infty,7}$.

The values of the $\delta_\square$ are given in Lemma 1 below and are achieved only with $\pm\gamma$ in $S_\square$. We should remark that the limit points are limit points of limit points from below.

We give the proof of Theorem 1 in §4 below.

The spectrum when a is odd.

Suppose now that a is odd. We suppose that

$$b = ma + r, \quad 0 < r \leq 2a, \quad 2 \mid r,$$

and set

$$n := m + 2, \quad s := m - 2,$$

so that m and n are the closest integers to b/a with the same parity as b.

We set

$$\begin{aligned}
S_0 &= \{\gamma : \ \gamma \text{ has periodic expansion, period } B_m\},\\
S_{-1} &= \{\gamma : \ \gamma \text{ has periodic expansion, period } B_m B_n\},\\
S_{-2} &= \{\gamma : \ \gamma \text{ has periodic expansion, period } B_n\},\\
S_{-3} &= \{\gamma : \ \gamma \text{ has periodic expansion, period } B_n B_s\},\\
S_{-4} &= \{\gamma : \ \gamma \text{ has periodic expansion, period } B_m B'_m \text{ (or } B_s B'_s \text{ if } m = 1)\},\\
S_{-5} &= \{\gamma : \ \gamma \text{ has periodic expansion, period } E_3 \ \},\\
S_{-6} &= \{\gamma : \ \gamma \text{ has periodic expansion, period } F_2 \ \},\\
S_{-7} &= \{\gamma : \ \gamma \text{ has periodic expansion, period } F_1 \text{ or } F_3 \ \},\\
S_{k,1} &= \{\gamma : \ \gamma \text{ has periodic expansion, period } B_n^k B_m\},\\
S_{k,2} &= \{\gamma : \ \gamma \text{ has periodic expansion, period } B_n^k B_m (B'_n)^k B'_m\},\\
S_{k,3} &= \{\gamma : \ \gamma \text{ has periodic expansion, period } (B_m B_n)^k B_n\},\\
S_{k,4} &= \{\gamma : \ \gamma \text{ has periodic expansion, period } (B_n B_m)^k (B'_n B'_m)^k\},\\
S_{k,5} &= \{\gamma : \ \gamma \text{ has periodic expansion, period } B_m^k B_n\},\\
S_{k,6} &= \{\gamma : \ \gamma \text{ has periodic expansion, period } (B_n B_s)^k B_n B_m^k\},\\
S_{k,7} &= \{\gamma : \ \gamma \text{ has periodic expansion, period } (B_n B_s)^k (B'_n B'_s)^k\},\\
S_{k,8} &= \{\gamma : \ \gamma \text{ has periodic expansion, period } (B_n B_s)^k B'_m\},\\
S_{k,9} &= \{\gamma : \ \gamma \text{ has periodic expansion, period } (B_m)^k B'_n E'_3 B'_s\},
\end{aligned}$$

and write

$$\delta_\square = M^*(\alpha, \gamma), \ \ \gamma \in S_\square.$$

In all cases except $a = 3, b = 4, 5, 6$ there are only finitely many values above the first limit point. In those cases (due to small partial quotients) the behaviour is atypical and we require additional sets of γ:

For $(a, b) = (3, 4)$ define

$$S_{-8} = \{\gamma : \ \gamma \text{ has periodic expansion, period } F_0B_0 \text{ or } F_2B_2' \ \},$$

$$S_{k,10} = \{\gamma : \ \gamma \text{ has periodic expansion, period } F_2(F_2B_2')^k \text{ or } F_2(F_0B_0)^k \ \}, \quad k \geq 1.$$

For $(a, b) = (3, 5)$ define blocks

$$G = \left(\frac{1}{2}(b-2-1), (a-1), \frac{1}{2}(b-2-1)\right),$$

$$H = \left(\frac{1}{2}(a-2+1), \frac{1}{2}(b-2-3), (a-1), \frac{1}{2}(b-2-3), \frac{1}{2}(a-2+1)\right),$$

$$H' = \left(\frac{1}{2}(a-2-1), \frac{1}{2}(b-2+1), (a-1), \frac{1}{2}(b-2+1), \frac{1}{2}(a-2-1)\right),$$

and sets

$$S_{-9} = \{\gamma : \ \gamma \text{ has periodic expansion, period } HGH'G \ \},$$

$$S_{k,11} = \{\gamma : \ \gamma \text{ has periodic expansion, period } (H'GHG)^kHG \ \}, \quad k \geq 0.$$

For $(a, b) = (3, 6)$ define

$$S_{k,12} = \{\gamma : \ \gamma \text{ has periodic expansion, period } F_2{B_2'}^{k+1} \text{ or } F_0B_2^kB_0 \ \}, \quad k \geq 0.$$

Theorem 2. *Suppose that $\alpha = [0; \overline{a, b}]$ with $a < b$ and $a \geq 3$ odd. Then*

$$\rho^*(\alpha) = \begin{cases} \delta_0, & \text{if } 2 \leq r \leq a-1, \ a \geq 5 \text{ or } a = 3, \ m \geq 2, \\ \delta_{-1}, & \text{if } r = a+1, \ a \geq 5 \text{ or } a = 3, \ m \geq 1, \\ \delta_{-2}, & \text{if } a+3 \leq r \leq 2a, \\ \delta_{-6}, & \text{if } (a, b) = (3, 4), \\ \delta_{-7}, & \text{if } (a, b) = (3, 5). \end{cases}$$

The remaining values $\rho_2(\alpha), \ldots$ down to the first limit point are as follows. In all cases $\delta_{k,i} \nearrow \delta_{\infty,i}$ as $k \to \infty$ for the appropriate $i = 1, \ldots, 9$. The values of these $\delta_{k,i}$ and δ_i are given in Lemma 2 below.

If $r \geq a+3$ then

$$\rho_2^*(\alpha) = \begin{cases} \delta_{\infty,1}, & \text{if } m \geq 1, \\ \delta_{\infty,2}, & \text{if } m = 0,\ a \geq 5. \end{cases}$$

If $r = a+1$ then

$$\rho_2^*(\alpha) = \begin{cases} \delta_{\infty,3}, & \text{if } m \geq 1, \\ \delta_{\infty,4}, & \text{if } m = 0,\ a \geq 5. \end{cases}$$

If $4 \leq r \leq a-1$ then

$$\rho_2^*(\alpha) = \begin{cases} \delta_{-5}, & \text{if } b = a+4 \geq 17, \\ \delta_{\infty,5}, & \text{otherwise,} \end{cases}$$
$$\rho_3^*(\alpha) = \delta_{\infty,5}, \quad \text{if } b = a+4 \geq 17.$$

If $r = 2$ then

$$\rho_2^*(\alpha) = \begin{cases} \delta_{-3}, & \text{if } m \geq 2, \\ \delta_{-5}, & \text{if } m = 1,\ a \geq 9, \end{cases}$$
$$\rho_3^*(\alpha) = \begin{cases} \delta_{\infty,6}, & \text{if } m \geq 3, \\ \delta_{\infty,7}, & \text{if } m = 2, \\ \delta_{\infty,9}, & \text{if } m = 1,\ a \geq 9. \end{cases}$$

For the remaining few cases:

If $(a,b) = (5,7)$ or $(7,9)$ then

$$\rho_2^*(\alpha) = \delta_{-3},$$
$$\rho_3^*(\alpha) = \delta_{-4},$$
$$\rho_4^*(\alpha) = \delta_{\infty,8}.$$

If $(a,b) = (3,4)$, then the values above $\delta_{\infty,10}$ are δ_{-6}, δ_{-8} and $\delta_{k,10} \searrow \delta_{\infty,10}$, $k \geq 0$.
If $(a,b) = (3,5)$ then the values above $\delta_{\infty,11}$ are δ_{-7}, δ_{-9} and $\delta_{k,11} \searrow \delta_{\infty,11}$, $k \geq 0$.
If $(a,b) = (3,6)$ then the values above $\delta_{\infty,12}$ are δ_{-2}, δ_{-6} and $\delta_{k,12} \searrow \delta_{\infty,12}$, $k \geq 0$.

We give the proof of Theorem 2 in §5 below.

THE SPECTRUM WHEN $a = 2$, $b \geq 5$.

We exclude here $b = 3, 4$ since the spectrum for $b = 3$ is the same as for the period one $[0; 3, \overline{4}]^- = 1 - [0; \overline{2, 3}]^-$ considered in [6] and the case $b = 4$, $[0; \overline{2, 4}] = 2 - \sqrt{2}$ has been considered already by Varnivides [7].

For $2 \leq t \leq b - 4$, with t of the same parity as b, define

$$S_{0,t} := \{\gamma \ : \ t_i \text{ periodic, period } a, -t \text{ or } a, t-4\}.$$

For b odd define

$$\begin{aligned}
S_{-2} &:= \{\gamma \ : \ t_i \text{ period } (a, -3, a, -1)\}, \\
S_{-1} &:= \{\gamma \ : \ t_i \text{ period } (a, -1)(a, -3, a, -1)(a, -1) \text{ or } (a, -3)(a, -1, a, -3)(a, -3)\}, \\
S_{2k+1} &:= \left\{\gamma \ : \ t_i \text{ period } (a, -1, 0, -1)(a, -3, a, -1)^k \text{ or } (a, -1, 0, -1)(a, -1, a, -3)^k\right\}, \quad k \geq 0, \\
S_{2k} &:= \left\{\gamma \ : \ t_i \text{ period } (a, -1)(a, -3, a, -1)^k \text{ or } (a, -3)(a, -1, a, -3)^k\right\}, \quad k \geq 1,
\end{aligned}$$

and for b even

$$\begin{aligned}
S_{-1} &:= \{\gamma \ : \ t_i \text{ period } 0, 0\}, \\
S_{2k+1} &:= \left\{\gamma \ : \ t_i \text{ period } (a, -2, 0, 0)(a, -2)^k \text{ or } (a, 0, 0, -2)(a, -2)^k\right\}, \quad k \geq 0, \\
S_{2k} &:= \left\{\gamma \ : \ t_i \text{ period } (a, -4)(a, -2)^k \text{ or } (a, 0)(a, -2)^k\right\}, \quad k \geq 1,
\end{aligned}$$

and set

$$\delta_k := M^*(\alpha, \gamma), \quad \gamma \in S_k.$$

Theorem 3. *Suppose that $\alpha = [0; \overline{2, b}]^-$ with $b \geq 5$.*

Then for b even

$$\rho^*(\alpha) = \begin{cases} \delta_{0,2} = \eta, & \text{if } b \text{ is even,} \\ \delta_{0,3} = \frac{2}{3}\eta, & \text{if } b = 5, \\ \delta_{0,3} = \eta\left(1 - \frac{1}{b^2 - b}\right), & \text{if } b \geq 7 \text{ is odd.} \end{cases}$$

The remaining values of $M(\alpha, \gamma)$ greater than

$$\delta_\infty = \eta\left((1-\beta)^2 - \begin{cases} \beta^2, & \text{if } b \text{ is even} \\ \frac{1}{4}\beta^4, & \text{if } b \text{ is odd.} \end{cases}\right),$$

are δ_{-1} (b even), δ_{-2}, δ_{-1} (b odd), the δ_{2k}, $\delta_{2k+1} \searrow \delta_\infty$, and when $b \geq 8$ the $\delta_{0,m}$ with $4 \leq m \leq 2 + [\sqrt{2b-4}]$ (m of the same parity as b). For these values $M^*(\alpha, \gamma) = \delta_j$ iff γ is in S_j.

3. The value of the points in the spectrum.

The following Lemmas give the values, δ_i, described in the above Theorems.

Lemma 1. *Suppose that $a \geq 4$ is even.*

When b is odd

$$\delta_0 = \left(1 - \beta - \frac{1}{b}\right)\left(1 - \eta + \frac{\eta}{b}\right),$$

$$\delta_{-1} = \left(1 - \beta - \frac{\beta(1-D)}{1+D^2}\right)\left(1 - \eta - \frac{D(1-D)}{1+D^2}\right),$$

$$\delta_{-2} = \begin{cases} \left(1 - \beta + \left(\frac{3\beta - 2D}{1-D}\right)\right)\left(1 - \eta - \left(\frac{2\eta - 3D}{1-D}\right)\right), & \text{if } b \geq \max\{2a-5, 3a/2\}, \\ \left(1 - \beta - \left(\frac{3\beta - 2D}{1-D}\right)\right)\left(1 - \eta + \left(\frac{2\eta - 3D}{1-D}\right)\right), & \text{if } b \leq \min\{a+5, 3a/2\}, \\ \left(1 + \beta - \left(\frac{3\beta - 2D}{1-D}\right)\right)\left(1 + \eta - \left(\frac{2\eta - 3D}{1-D}\right)\right), & \text{otherwise,} \end{cases}$$

and

$$\delta_k = \left(1 - \beta - \frac{\beta(1-D)(1+2D^2)}{1+D^2} - \beta D^3 w_k\right)\left(1 - \eta - \frac{D(1-D)}{1+D^2} + w_k\right)$$

where $w_k := 2D^{4k+1}(1-D)/((1+D^2)(1-D^{4k+2}))$.

When b is even,

$$\delta_{k,1} = \left(1 - \eta + \frac{2D^2}{1+D}(1 - w_{k,1})\right)\left(1 - \beta - \frac{2\beta}{1+D}(1 - w_{k,1})\right)$$

where $w_{k,1} := D^{2k}(1-D)/(1-D^{2k+1})$, $k \geq 0$.

For $b = 2a - 2$, $a \geq 8$,

$$\delta_{k,2} = \left(1 - 3\eta + \frac{2D(2-\eta)}{1-D} - w_{k,2}\right)\left(1 + \beta - \frac{2\beta D}{1-D}(1 - \eta + D) + \beta D w_{k,2}\right)$$

where $w_{k,2} := 2D^{k+1}(1+D)(1-\eta+D)/((1-D)(1+D^{k+2}))$.

For $b = 2a - 4$, $a \geq 10$,

$$\delta_{k,3} = \left(1 - \eta - \frac{2\eta D}{1-D} + \frac{2D}{1-D^2}(1+2D) + w_{k,3}\right) \left(1 - 3\beta + \frac{2D}{1-D} - \frac{2\beta D}{1-D^2}(2+D) - \beta D w_{k,3}\right)$$

with $w_{k,3} := 2D^{2k+1}/((1+D)(1-D^{2k+1}))$.

For $a + 6 \leq b \leq 2a - 6$

$$\delta_{k,4} = \left(1 - \eta + \frac{2D(1-\eta)}{1-D} + w_{k,4}\right)\left(1 - 3\beta + \frac{2D(1-\beta)}{1-D} - \beta w_{k,4}\right)$$

where $w_{k,4} := 2D^{k+1}/(1-D^{k+1})$.

For $k = 0$ *or* $b \leq 2a - 6$ *or* $a = 6, b = 8$, *or* $k = 1$ *and* $b = 2a - 4$

$$\delta_{k,5} = \left(1 - \eta + \frac{2D(1-\eta)}{1-D} + \eta w_{k,5}\right)\left(1 - 3\beta + \frac{2D(1-\beta)}{1-D} - w_{k,5}\right)$$

where $w_{k,5} := 2D^{k+1}(1 - 2\beta + D)/((1-D)(1+D^{k+1}))$.

For $(a, b) = (8, 12)$

$$\delta_{k,6} = \left(1 - 3\eta + 2D - 2D^2 + 2\eta D^2 - \frac{2D^3(1-\eta+D)}{1+D^2} - \eta D^2 w_{k,6}\right) \left(1 + \beta - \frac{2D(1-\beta+\beta D)}{1+D^2} + w_{k,6}\right)$$

where $w_{k,6} := 2D^{4k+4}(1 - 2\beta + D)(1 - D^3)/((1 + D^2)(1 - D^{4k+6}))$.

For $(a, b) = (6, 10)$

$$\delta_{k,7} = \left(1 + \eta - 2D + 2D^2 + \frac{2\eta D^3}{1-D} - \frac{2D^3(1+2D)}{1-D^2} - 2\eta D^3 w_{k,7}\right) \left(1 - 5\beta + \frac{2D}{1-D} - 2\beta D\frac{(1+2D)}{(1-D^2)} - w_{k,7}\right)$$

where $w_{k,7} := 2D^{2k}(1 - 3\beta + D)/(1 - D^{2k+2})$, *and* $\delta_{1,4} = \frac{703}{40} - \frac{703}{600}\sqrt{210}$.

For the remaining δ_i *we have*

$$\delta_{-1} = \left(1 - 3\eta + \frac{2D(1-\eta)}{1-D}\right)\left(1 - \beta + \frac{2\beta(1-\eta)}{1-D}\right)$$

and

$$\delta_{0,4} = \begin{cases} \left(1+3\beta-\frac{2D(1-2\beta)}{1-D}\right)\left(1-3\eta+\frac{2D(2-\eta)}{1-D}\right), & \text{if } b \geq \max\{3a-6, 2a\}, \\ \left(1-5\beta+\frac{2D(1-2\beta)}{1-D}\right)\left(1+\eta-\frac{2D(2-\eta)}{1-D}\right), & \text{if } b \leq \min\{a+6, 2a-2\}, \\ \left(1-3\beta+\frac{2D(1-2\beta)}{1-D}\right)\left(1-\eta+\frac{2D(2-\eta)}{1-D}\right), & \text{else,} \end{cases}$$

and for $(a,b) = (4,6)$

$$\delta_{-2} = \left(1-\eta-\frac{2D}{1-D}+\frac{2\eta D}{1-D^2}\right)\left(1-3\beta-\frac{2\beta D}{1-D}+\frac{2D}{1-D^2}\right).$$

Lemma 2. *Suppose that* a *is odd.*

Set

$$v := \frac{m\beta - D}{1-D}, \quad D := \eta\beta.$$

Then

$$\delta_0 = (1-2\eta+\eta v)(1-\beta+v), \tag{3.1}$$

$$\delta_{-1} = \begin{cases} \left(1-\eta v-\frac{2D^2}{1-D^2}\right)\left(1-3\beta-v-\frac{2\beta D^2}{1-D^2}\right), & \text{if } r \leq a+1, \\ \left(1-2\eta+\eta v+\frac{2D}{1-D^2}\right)\left(1-\beta+v+\frac{2\beta D}{1-D^2}\right), & \text{if } r \geq a+1, \end{cases} \tag{3.2}$$

$$\delta_{-2} = \left(1-\eta v-\frac{2D}{1-D}\right)\left(1-3\beta-v-\frac{2\beta D}{1-D}\right), \tag{3.3}$$

$$\delta_{-3} = \left(1-2\eta+\eta v+\frac{2\eta}{b}\right)\left(1-3\beta+v+\frac{2D}{b}\right), \tag{3.4}$$

$$\delta_{-4} = \left(1-2\eta+\frac{\eta(m+\eta)}{b}\right)\left(1-\beta-\frac{(m-\eta)}{b}\right). \tag{3.5}$$

If $m = 1$ *then*

$$\delta_{-5} = \begin{cases} \left(1-4\eta+\frac{3D(1-\eta)}{1-D}\right)\left(1+2\beta-\frac{3D(1-\beta)}{1-D}\right), & \text{if } r \geq a-7, \\ \left(1-2\eta+\frac{3D(1-\eta)}{1-D}\right)\left(1-2\beta+\frac{3D(1-\beta)}{1-D}\right), & \text{if } r \leq a-9. \end{cases} \tag{3.6}$$

For b even

$$\delta_{-6} = \eta$$

and for b odd

$$\delta_{-7} = \eta\left(1 - \left(\frac{\beta}{1-D}\right)^2\right).$$

For $r \geq a+3$

$$\delta_{k,1} = \left(1 - 2\eta + \eta v + \frac{2D}{1-D} - \frac{2D^{k+1}}{1-D^{k+1}}\right)\left(1 - \beta + v + \frac{2\beta D}{1-D} - \frac{2\beta D^{k+1}}{1-D^{k+1}}\right). \tag{3.7}$$

For $m = 0$, $k \geq 1$, $r \geq a+3$,

$$\delta_{k,2} = \left(1 - 2\eta + \frac{D(2-\eta)}{1-D} - \eta\varepsilon_{k,2}\right)\left(1 - \beta + \frac{D(1-2\beta)}{1-D} + D\varepsilon_{k,2}\right). \tag{3.8}$$

For $r \leq a+1$ and $b \geq 6$

$$\delta_{k,3} = \left(1 - \eta v - \frac{2D}{1-D^2} - \eta\varepsilon_{k,3}\right)\left(1 - 3\beta - v - \frac{2\beta D^2}{1-D^2} - \varepsilon_{k,3}\right). \tag{3.9}$$

For $b = a+1 \geq 6$

$$\delta_{k,4} = \left(1 - \frac{\eta D}{1-D} + \frac{2D^2}{1-D^2} + \eta D\varepsilon_{k,4}\right)\left(1 - 3\beta + \frac{D}{1-D} - \frac{2\beta D^2}{1-D^2} - \varepsilon_{k,4}\right). \tag{3.10}$$

For $r \leq a-1$

$$\delta_{k,5} = \left(1 - \eta v - \frac{2D^{k+1}}{1-D^{k+1}}\right)\left(1 - 3\beta - v - \frac{2\beta D^{k+1}}{1-D^{k+1}}\right). \tag{3.11}$$

For $r = 2$ and $b \geq 7$

$$\delta_{k,6} = \left(1 - \eta v - \frac{2D^{k+1}(1+D^{2k+1})}{(1+D)(1-D^{3k+1})}\right)\left(1 - 3\beta - v + \frac{2D}{b} - \frac{2\beta D^{2k+1}(1+D^k)}{(1+D)(1-D^{3k+1})}\right). \tag{3.12}$$

For $b = 2a+2$

$$\delta_{k,7} = \left(1 - \frac{\eta D}{1-D} + \frac{4D^2}{1-D^2} + \eta D\varepsilon_{k,7}\right)\left(1 - \beta + \frac{D}{1-D} - \frac{4\beta}{1-D^2} - \varepsilon_{k,7}\right). \tag{3.13}$$

For $b = a + 2 \geq 7$

$$\delta_{k,8} = \left(1 + \frac{D(1+D-4D^2)}{1-D^2} - \frac{\eta D(1-2D)}{1-D} - \eta D^2 \varepsilon_{k,8}\right) \left(1 - 4\beta + \frac{D}{1-D} + \frac{\beta D(1-3D)}{1-D^2} - \varepsilon_{k,8}\right). \tag{3.14}$$

and $b = a + 2 \geq 11$

$$\delta_{k,9} = \left(1 - 2\eta + 3D - 3\eta D + 3D^2 - \eta D^2 - \frac{\eta D^2(\beta - D)}{1-D} - \eta D^2 \varepsilon_{k,9}\right) \left(1 - 2\beta + \frac{D(1-\beta)}{1-D} - \varepsilon_{k,9}\right). \tag{3.15}$$

Here the $\varepsilon_{k,i} \to 0$ *are given explicitly by*

$$\varepsilon_{k,2} := \frac{2D^k\,(\beta(1+D) - D)}{(1-D)(1+D^{k+1})}, \quad \varepsilon_{k,3} := \frac{2\beta D^{2k+1}}{(1+D)(1-D^{2k+1})},$$

$$\varepsilon_{k,4} := \frac{2D^{2k}(1-2/b)}{(1-D)(1+D^{2k})}, \quad \varepsilon_{k,7} := \frac{2D^{2k}(1-4/b)}{(1-D)(1+D^{2k})},$$

$$\varepsilon_{k,8} := \frac{2D^{2k}(1-2/b)}{1-D^{2k+1}}, \quad \varepsilon_{k,9} := \frac{2D^{k+1}}{1-D^{k+3}}(1 - 2\beta + 2D - 2\beta D + D^2).$$

For $(a, b) = (3, 4)$

$$\delta_{-8} = \eta\left(1 - \left(\frac{\beta(1-\eta+D)}{1-D^2}\right)^2\right),$$

$$\delta_{k,10} = \eta\left(1 - \frac{\beta(1-\eta+D)}{1-D^2} + \varepsilon_{k,10}\right)\left(1 + \frac{\beta D(1-\eta+D)}{1-D^2} - D\varepsilon_{k,10}\right)$$

where $\varepsilon_{k,10} = \beta D^{2k}(1-\eta+D)/(1+D)(1-D^{2k+1})$.

For $(a, b) = (3, 5)$

$$\delta_{-9} = \eta\left(1 - \left(\frac{2\beta - D + D^3 - 2\beta D^3}{1+D^4}\right)^2\right),$$

$$\delta_{k,11} = \eta\left(1 - 2\beta + D - D^3\frac{(1 - 2\beta - 2\beta D + D^2)}{1+D^4}(1 - \varepsilon_{k,11})\right) \left(1 + 2\beta - D - D^3\frac{(1 - 2\beta - 2\beta D + D^2)}{1+D^4}(1 + D^4\varepsilon_{k,11})\right)$$

where $\varepsilon_{k,11} = 2D^{8k}/(1 - D^{8k+4})$.

For $(a, b) = (3, 6)$

$$\delta_{k,12} = \eta\left(1 - \left(\frac{\beta(1-\eta+D)(1-D^{k+1})}{(1-D)(1-D^{k+2})}\right)^2\right).$$

Lemma 3. *Suppose that $a = 2$. For $2 \le t \le b$*

$$\delta_{0,t} = \begin{cases} \eta\left(1 - \left(\frac{(t-2)\beta}{1-D}\right)^2\right), & \text{if } t \le b - \sqrt{2b-4}, \\ \eta\left(\left(2 - 2\beta - \frac{(t-2)\beta}{1-D}\right)^2 - 1\right), & \text{if } t > b - \sqrt{2b-4}. \end{cases}$$

For b even

$$\begin{aligned} \delta_{-1} &= \eta(1-\beta)^2, \\ \delta_{2k} &= \eta\left(1 - 2\beta - \frac{2\beta D^{k+1}}{1-D^{k+1}}\right)\left(1 + \frac{2\beta D^k}{1-D^{k+1}}\right), \\ \delta_{2k+1} &= \eta\left((1-\beta)^2 - \beta^2\left(\frac{1-D^{k+1}}{1-D^{k+2}}\right)^2\right). \end{aligned}$$

For b odd

$$\begin{aligned} \delta_{-2} &= \eta\left(1 - \beta + \frac{\beta D}{1+D}\right)^2, \\ \delta_{-1} &= \eta\left(\left(1 - \beta + \frac{\beta(1-D)D^3}{1-D^4}\right)^2 - \left(\frac{\beta(1+D)D}{1-D^4}\right)^2\right), \\ \delta_{2k} &= \eta\left(1 - \beta - \frac{1}{2}\beta^2 - \frac{2\beta D^{2k+2}}{(1+D)(1-D^{2k+1})}\right)\left(1 - \beta + \frac{1}{2}\beta^2 + \frac{2\beta D^{2k}}{(1+D)(1-D^{2k+1})}\right), \\ \delta_{2k+1} &= \eta\left((1-\beta)^2 - \frac{1}{4}\beta^4\left(\frac{1-D^{2k}}{1-D^{2k+2}}\right)^2\right). \end{aligned}$$

Proof of Lemma 1 (even $a \ge 4$)

Notice that (because of the forwards backwards symmetry, and the rough bound $s_4^*(2i-1) \ge (1+\eta-D)(1+\beta) > 1$ when $d_{2i-1}^+ > 0$) in all cases it is enough to check

$$u_i := \min\{s_1^*(2i-1), s_2^*(2i-1), s_3^*(2i-1)\}, \qquad d_{2i-1}^+ \ge 0,$$

for each i in the period of γ or its negative, where $s_1^*(2i-1) < s_2^*(2i-1)$ iff $(d_{2i-1}^+ - \beta)(1 + d_{2i-1}^-) < \eta$, with $s_1^*(2i-1) < s_3^*(2i-1)$ iff $d_{2i-1}^-(1-\beta) + d_{2i-1}^+(1-\eta) < 0$ and $s_2^*(2i-1) < s_3^*(2i-1)$ iff $(d_{2i-1}^- + \eta)(1 - d_{2i-1}^+) + \beta < 0$.

For period C_3 we have $d^+_{2i-1} = (3\beta - 2D)/(1-D)$, $d^-_{2i-1} \sim (-2\eta + 3D)/(1-D)$ and one tediously checks which of the three functions gives u_i.

When $(b_{2i-1}, b_{2i}) = A_0$, A_1, A_2, or C_2 the bound $d^+_{2i-1} \le 2\beta$ plainly gives $s^*_1(2i-1) < s^*_2(2i-1)$.

For blocks C_2 the bounds $d^-_{2i-1} \le -2\eta + 4D$, $d^+_{2i-1} \le 2\beta$ give $s^*_1(2i-1) < s^*_3(2i-1)$ and hence $u_i = s^*_1(2i-1)$.

For blocks A_1 or A_2 the bounds $d^-_{2i-1} \ge -2D$, $d^+_{2i-1} \ge \beta - D$ give $u_i = s^*_3(2i-1)$.

Hence the values of $\delta_{0,1}$, $\delta_{0,5}$ and δ_{-1} with b even, and δ_0 when b is odd.

Observe that if a block A_1 is preceded by an A'_1 then $d^+_{2i-1} \le \beta + \beta D$, $d^-_{2i-1} \le -D + 2D^2$ gives $s^*_3(2i-1) \ge (1 - 2\beta - \beta D)(1 - \eta + D - 2D^2) > (1-2\beta)(1-\eta)$ greater than the value claimed for δ_{-1}, b odd and the δ_k, $k \ge 1$. Hence in these cases we need only check the A_1 preceded by an A_1. For δ_{-1} these have $d^+_{2i-1} = \beta(1-D)/(1+D^2)$, $d^-_{2i-1} \sim D(1-D)/(1+D^2)$, and $s^*_3(2i-1)$ gives δ_{-1}. For blocks $A'_1A_1(A'_1A'_1A_1A_1)^k$ observe that taking the negative essentially reverses the sequence (interchanging d^-_{2i-2} and d^+_{2i-1}) and we need just check $s_3 := (1 - \eta - \eta d)(1 - \beta - d')$ where $d = \min\{d^-_{2i-2}, d^+_{2i-1}\}$, $d' = \max\{d^-_{2i-2}, d^+_{2i-1}\}$ for $t_{2i} = 1 = t_{2i-2}$. Plainly the largest d, d' both occur when the A_1 is followed by $A'_1A_1A'_1$ and

$$d' \sim \frac{\beta(1-D)}{1-D^{4k+2}}\left(1 + \frac{D^2(1-D^{4k})}{1+D^2}\right), \quad d \sim \frac{\beta(1-D)}{1-D^{4k+2}}\left(\frac{(1-D^{4k})}{1+D^2} - D^{4k}\right),$$

with $s^*_3(2i-1)$ giving the value claimed for δ_k, $k \ge 1$.

For the blocks A_0 in $S_{k,1}$, $k \ge 1$, the bounds $d^+_{2i-1} \le 2\beta D$, $d^-_{2i-1} \le -2D + 2D^2$ give $u_i = s^*_1(2i-1) \sim \left(1 - \eta - \frac{2D}{1+D}(1 - w_{k,1})\right)\left(1 - \beta + \frac{2\beta D}{1+D}(1 - w_{k,1})\right)$, while cutting at the A_2 the minimum value of the $u_i = s^*_3(2i-1)$ occurs when $d^+ = 2\beta(1-w_{k,1})/(1+D)$, $d^- \sim -2D^2(1-w_{k,1})/(1+D)$ (i.e. when d^- and d^+ are both largest) with this giving the value claimed for $\delta_{k,1}$.

For the $\delta_{k,5}$ observe that cutting at an A_2 gives

$$d^- \sim \left(-2D(1-\eta)\frac{(1-D^k)}{(1-D)} - 2D^{k+1}\right)/(1+D^{k+1}),$$

$$d^+ = \left(2\beta - 2D(1-\beta)\frac{(1-D^k)}{(1-D)}\right)/(1+D^{k+1})$$

and $u_i = s_3^*(2i-1)$ asymptotically gives the value claimed for $\delta_{k,5}$ with these increasing to $\delta_{\infty,5} = \delta_{\infty,4}$. Cutting at a C_2 gives $d^- \geq (-2\eta + 2D)/(1-D)$ and $d^+ \geq (2\beta - 2D)/(1-D)$ and

$$\begin{aligned} u_i = s_1^*(2i-1) &\geq \left(1 - \eta - \frac{2\eta}{1-D}(1-\beta)\right)\left(1 - \beta + \frac{2\beta}{1-D}(1-\eta)\right) \\ &= \delta_{\infty,5} + \frac{2D}{(1-D)^2}(2a - 5 - b + 7\eta - 4D - \eta D + D^2) > \delta_{\infty,5} \end{aligned}$$

for $b \leq 2a-6$ or $a = 6, b = 8$. For $k = 1$ and $b = 2a-4$ one checks that these

$$u_i = s_1^*(2i-1) \sim \left(1 - 3\eta + 2D\frac{(1 - D + \eta D)}{1+D^2}\right)\left(1 + \beta - 2\beta D\frac{(1 - \eta + D)}{1+D^2}\right)$$

are still greater than the value claimed for $\delta_{1,5}$;

$$u_i = s_3^*(2i-1) \sim \left(1 - \eta + 2D\frac{(1 - \eta + D)}{1+D^2}\right)\left(1 - 3\beta + 2D\frac{(1 - \beta + \beta D)}{1+D^2}\right).$$

Cutting at a C_4 the bounds

$$\frac{(4\beta - 2D)}{(1-D)} \geq d^+ \geq (4\beta - 2D), \quad \frac{(-2\eta + 4D)}{(1-D)} \geq d^- \geq -2\eta + 2D - 2\eta D,$$

give $s_1^*(2i-1) \leq s_3^*(2i-1)$ iff $b \geq 2a$, give $s_2^*(2i-1) < s_3^*(2i-1)$ if $b \geq a+8$ (or $b = a+6$ when C_4 is not preceded by another C_4, using $d^- \leq -2\eta + 2D$) and $s_2^*(2i-1) > s_3^*(2i-1)$ if $b \leq a+4$ (or $b = a+6$ when C_4 is preceded by another C_4, using $d^- \geq -2\eta + 4D - 2\eta D$), and give $s_1^*(2i-1) < s_2^*(2i-1)$ if $b \geq \max\{3a-6, 2a\}$ and $s_2^*(2i-1) < s_1^*(2i-1)$ if $b \leq 3a-8$. Hence the expressions for $\delta_{0,4}$.

For the $S_{k,2}$ observe that cutting at a C_2 produces

$$\begin{aligned} d^+ &= \left(2\beta - 2\beta D + 2D^2(1-2\beta)\frac{(1-D^k)}{(1-D)} + 2D^{k+2}\right)/(1+D^{k+2}) \\ d^- &\sim \left(-2\eta + 2D(2-\eta)\frac{(1-D^k)}{(1-D)} + (2-2D)D^{1+k}\right)/(1+D^{k+2}) \end{aligned}$$

and $u_i = s_1^*(2i-1)$ asymptotically gives the value claimed for $\delta_{k,2}$ with these increasing to $\delta_{\infty,2} \leq 1 - 3\eta + \beta + D - 2\eta D + 2\beta D + 10D^2$. Cutting at an A_2 the bounds $d^- \leq -2D + 2\eta D$, $d^+ \leq 2\beta - 2D + 4\beta D$, give

$$\begin{aligned} u_i = s_3^*(2i-1) &\geq (1 - \eta + 2D - 2\eta D)(1 - 3\beta + 2D - 4\beta D) \\ &\geq 1 - \eta - 3\beta + 7D - 4\eta D - 10\beta D + 10D^2 \\ &\geq \delta_{\infty,2} + 2D(b - 2a + 3 - 2\eta - 4\beta) > \delta_{\infty,2} \end{aligned}$$

for $b = 2a-2$, $a \geq 6$. Cutting at a C_4 the bounds $d^+ \leq 4\beta - 2D + 4\beta D$, $-2\eta + 4d \geq d^- \geq -2\eta + 2D - 2D^2$, give

$$\begin{aligned} s_2^*(2i-1) &\geq (1-\eta+2D-2D^2)(1-3\beta+2D-4\beta D) \\ &\geq 1-3\beta-\eta+7D-2\eta D-10\beta D \geq \delta_{\infty,2} + 2D(b-2a+3-\eta-4\beta-5D) \end{aligned}$$

(and $s_3^*(2i-1) > \delta_{\infty,2}$ for $(a,b) = (8,14)$). Hence the value of $\delta_{k,2}$ for $b = 2a-2$, $a \geq 8$.

For the $\delta_{k,3}$ observe that for the C_4 the smallest d^- and largest d^+ both occur when

$$d^+ = \frac{4\beta - 2D}{1-D} - \frac{2\beta D^2(1-D^{2k})}{(1-D^2)(1-D^{2k+1})}, \quad d^- \sim \frac{-2\eta + 4D}{1-D} - \frac{2D(1-D^{2k})}{(1-D^2)(1-D^{2k+1})},$$

so that when $a+6 \leq b < 2a$ we have $u_i = s_2^*(2i-1) \leq (1-\eta+2D)(1-3\beta+2D)$ giving the value claimed for $\delta_{k,3}$ (checking numerically that for $(a,b) = (10,16)$ one has $u_i = s_3^*(2i-1) \geq (1+\eta-4D+2\eta D-2D^2)(1-5\beta+2D-2\beta D)$ for a C_4 preceded by a C_4, greater than the claimed value $\delta_{k,3}$). Cutting at a C_2 the rough bounds $d^- \geq -2\eta+4D-2\eta D$, $d^+ \geq 2\beta - 2D$ give $u_i = s_1^*(2i-1) \geq (1-3\eta+4D-2\eta D)(1+\beta-2D)$, which is larger when $b = 2a-4$.

For the $\delta_{k,4}$ cutting at the C_4 gives

$$d^+ = \frac{2\beta - 2D}{1-D} + \frac{2\beta}{1-D^{k+1}}, \quad d^- \sim \frac{-2\eta + 2D}{1-D} + \frac{2D^{k+1}}{1-D^{k+1}}$$

and when $a+6 \leq b < 2a$

$$u_i = s_2^*(2i-1) \sim \left(1-\eta+\frac{2D(1-\eta)}{1-D} + w_{k,4}\right)\left(1-3\beta+\frac{2D(1-\beta)}{1-D} - \beta w_{k,4}\right)$$

gives the value claimed. Cutting at a C_2

$$\begin{aligned} u_i = s_1^*(2i-1) &\geq \left(1-3\eta+\frac{2D-2\eta D}{1-D}\right)\left(1+\beta-\frac{(2D-2\beta D)}{1-D}\right) \\ &\geq \left(1-\eta+\frac{4D-2\eta D}{1-D}\right)\left(1-3\beta+\frac{2D-4\beta D}{1-D}\right) \end{aligned}$$

(the value claimed when $k=0$) for $b \leq 2a-6$.

For $(a,b) = (8,12)$ and $\delta_{k,6}$ cutting at the first C_2 in a block C_2C_2 gives

$$d^- \sim \left(-2\eta + 2D + 2\eta D^2 - 2D^2 - 2D^3\frac{(1-\eta+D)(1-D^{4k+4})}{1+D^2}\right)/(1-D^{4k+6}),$$

$$d^+ \sim \left(2\beta \frac{(1-\eta+D)(1-D^{4k+4})}{1+D^2} + 2D^{4k+4}(1-\beta+\beta D-D^2) \right) / (1-D^{4k+6})$$

and $u_i = s_1^*(2i-1)$ asymptotically gives the value claimed for $\delta_{k,6}$ (the largest value corresponding to $k=0$). For the remaining C_2 and A_2 we have the even larger

$$u_i = s_1^*(2i-1) \geq (1-3\eta+2D-2\eta D)(1+\beta-2\beta D)$$

and

$$u_i = s_3^*(2i-1) \geq (1-\eta+2D-2\eta D)(1-3\beta+2D-2\beta D).$$

For $(a,b)=(6,10)$ cutting at the C_4 in blocks A_2C_4 gives $s_3^*(2i-1) \leq \delta_{\infty,7}$ asymptotically equal to the value claimed for $\delta_{k,7}$. For the remaining C_4

$$u_i = s_3^*(2i-1) \geq (1-5\beta+2D-2\beta D)(1+\eta-2D+2\eta D-4D^2) > \delta_{\infty,7}$$

and for the A_2 and C_2 respectively we have

$$u_i = s_3^*(2i-1) \geq (1-\eta+2D-2D^2)(1-3\beta+2D-4\beta D+2D^2-2\beta D^2) > \delta_{\infty,7},$$

$$u_i = s_1^*(2i-1) \geq \left(1+\beta-2D+4\beta D-2D^2\right)(1-3\eta+4D-2\eta D) > \delta_{\infty,7}.$$

Finally when $(a,b)=(4,6)$, in the case of δ_{-2}, cutting at an A_2 we have $u_i = s_3^*(2i-1)$ asymptotically giving the value claimed, while cutting at the C_2 gives

$$u_i = s_1^*(2i-1) \sim \left(1-3\eta+\frac{2D}{1-D}-\frac{2\eta D^2}{1-D^2}\right)\left(1+\beta+\frac{2\beta D}{1-D}-\frac{2D^2}{1-D^2}\right). \quad \blacksquare$$

Proof of Lemma 2 (odd a)

We first exclude the S_k containing $t_{2i-1} = a$. Hence it is enough to check the minimum of the $s_j^*(2i), s_j^*(2i+1)$ for $j = 1, ..., 4$ for all i in the period of $\pm\gamma$ with $t_{2i+1} < 0$. Since the $d_{2i}^-, d_{2i+1}^+ \geq -2\beta - D$ we can ignore the $s_4^*(2i+1), s_2^*(2i) \geq (1+\eta)(1-\beta-D) > (1-\eta)(1-\beta)$ and need only check

$$v_i := \min\{s_1, s_3, s_2\}, \qquad t_{2i+1} < 0,$$

where, writing $d := \min\{d^+_{2i+1}, d^-_{2i}\}$, $d' := \max\{d^+_{2i+1}, d^-_{2i}\}$,

$$s_1 := \min\{s^*_1(2i+1), s^*_1(2i)\} = (1-\eta+t_{2i+1}\eta+\eta d')(1-\beta+d),$$
$$s_3 := \min\{s^*_3(2i+1), s^*_3(2i)\} = (1-\eta-t_{2i+1}\eta-\eta d)(1-\beta-d'),$$
$$s_2 := \min\{s^*_2(2i+1), s^*_4(2i)\} = (1+\eta+t_{2i+1}\eta+\eta d)(1+\beta-d').$$

Now if $t_{2i+1} = -1$ then $v_i = \min\{s_1, s_3\}$ (if $d' > 0$ then $d \geq -2\beta - D$ gives $-\eta d(1-d') < \beta$ and $s_3 < s_2$, while if $d' < 0$ then plainly $s_1 < s_2$). For γ in S_{-4} we have $d = -(m\beta - D)/(1+D)$, $d' = (m\beta+D)/(1+D)$ and $v_i = s_1$ gives the value of δ_{-4}.

We first deal with the γ having $t_{2i+1} = -1$ for all i. Writing

$$d = \frac{m\beta - D}{1-D} + 2\beta w, \quad d' = \frac{m\beta - D}{1-D} + 2\beta w',$$

for $t_{2i+1} = -1$ we have

$$s_1 < s_3 \text{ iff } w(1-\eta-D) + w'(1+\eta-D) < \frac{r}{a}.$$

In particular

$$\{t_{2i}, t_{2i+2}\} = \{m, m\} \Rightarrow v_i = s_1$$

giving (3.1) for δ_0. Likewise

$$\{t_{2i}, t_{2i+2}\} = \{n, n\} \Rightarrow v_i = \begin{cases} s_1, & \text{if } r = 2a, \\ s_3, & \text{if } r \leq 2a-2, \end{cases}$$

(with asymptotic equality when $r = 2a$ and $t_{2i} = n$ for all i) and (3.3) is clear. Similarly

$$\{t_{2i}, t_{2i+2}\} = \{n, m\} \Rightarrow v_i = \begin{cases} s_1, & \text{if } r \geq a+3, \\ s_3, & \text{if } r \leq a-1, \end{cases}$$

where when $r = a+1$ we have $v_i = s_3$ if $w' \geq 1/(1-D^2)$, $w \geq D/(1-D^2)$ and $v_i = s_1$ if $w' \leq 1/(1-D^2)$, $w \leq D/(1-D^2)$ (with asymptotic equality when the t_{2i} have period n, m) giving the value (3.2) of δ_{-1}. For the γ in S_{-3} we have $w' = 1/(1+D)$, $w = -1/(1+D)$ and s_1 gives δ_{-3}.

Suppose $r \geq a+3$ and γ is in $S_{k,1}$. If $\{t_{2i}, t_{2i+2}\} = \{m,n\}$ then $d' = v + 2\beta/(1-D) - 2\beta D^k/(1-D^{k+1})$ and $d = v + 2\beta/(1-D) - 2\beta/(1-D^{k+1})$ and

$$\begin{aligned} v_i = s_1 &= \left(1 - 2\eta + \eta v + \frac{2D}{1-D} - \frac{2D^{k+1}}{1-D^{k+1}}\right)\left(1 - \beta + v + \frac{2\beta D}{1-D} - \frac{2\beta D^{k+1}}{1-D^{k+1}}\right) \\ &< \delta_{\infty,1} := \left(1 - 2\eta + \eta v + \frac{2D}{1-D}\right)\left(1 - \beta + v + \frac{2\beta D}{1-D}\right). \end{aligned}$$

For the remaining $\{t_{2i}, t_{2i+2}\} = \{n,n\}$ we have $v + 2\beta < d, d' < v + 2\beta/(1-D)$ and

$$\begin{aligned} s_1 &\geq (1 - 2\eta + \eta v + 2D)(1 - \beta + v + 2\beta) \\ &> \delta_{\infty,1} + \frac{2\beta}{1-D}(1-2D)(1-2\eta) - \frac{2D^2}{1-D}(1+\eta) > \delta_{\infty,1}, \end{aligned}$$

while

$$\begin{aligned} s_3 &\geq \left(1 - \eta v - \frac{2D}{1-D}\right)\left(1 - \beta - v - \frac{2\beta}{1-D}\right) \\ &= \delta_{\infty,1} + 2\beta\left(\frac{r-(a+3)}{a} + \eta\left(1 - \frac{3\beta}{a} + \frac{\beta(m+2) - D}{1-D}\right)\right) > \delta_{\infty,1}, \end{aligned}$$

and (3.7) holds.

Suppose that $r \leq a-1$ and γ is in $S_{k,5}$. If $\{t_{2i}, t_{2i+2}\} = \{m,n\}$ we have $d' = v + 2\beta/(1-D^{k+1})$, $d = v + 2\beta D^k/(1-D^{k+1})$ and

$$\begin{aligned} v_i = s_3 &= \left(1 - \eta v - \frac{2D^{k+1}}{1-D^{k+1}}\right)\left(1 - 3\beta - v - \frac{2\beta D^{k+1}}{1-D^{k+1}}\right) \\ &< \delta_{\infty,5} := (1 - \eta v)(1 - 3\beta - v). \end{aligned}$$

For the remaining $\{t_{2i}, t_{2i+2}\} = \{m,m\}$ we have $d, d' \geq v$ and

$$\begin{aligned} v_i = s_1 &\geq (1 - 2\eta + \eta v)(1 - \beta + v) \\ &= \delta_{\infty,5} + 2\beta\left(\frac{a-1-r}{a} + \frac{1}{a} - \eta v\right) > \delta_{\infty,5}, \end{aligned}$$

giving (3.11).

Suppose that $r \leq a+1$ and γ is in $S_{k,3}$. If $\{t_{2i}, t_{2i+2}\} = \{n,n\}$ we have $d, d' = v + 2\beta/(1-D^2) + \varepsilon_{k,3}$ and

$$\begin{aligned} v_i = s_3 &= \left(1 - \eta v - \frac{2D}{1-D^2} - \eta\varepsilon_{k,3}\right)\left(1 - 3\beta - v - \frac{2\beta D^2}{1-D^2} - \varepsilon_{k,3}\right) \\ &< \delta_{\infty,3} := \left(1 - \eta v - \frac{2D}{1-D^2}\right)\left(1 - 3\beta - v - \frac{2\beta D^2}{1-D^2}\right). \end{aligned}$$

For the other $\{t_{2i}, t_{2i+2}\} = \{n, m\}$ we have $d' \le v+2\beta/(1-D)$ and $d \le v+2\beta D/(1-D)$ and for $b \neq 4, 5$

$$\begin{aligned} v_i = s_3 &\ge \left(1 - \eta v - \frac{2D^2}{1-D}\right)\left(1 - \beta - v - \frac{2\beta}{1-D}\right) \\ &\ge \delta_{\infty,3} + \frac{2D}{1-D^2}(1 - (4+m)\beta) > \delta_{\infty,3}, \end{aligned}$$

giving (3.9) for $b \neq 4, 5$.

Suppose that $r = 2$ and that γ is in $S_{k,6}$. If $\{t_{2i}, t_{2i+2}\} = \{n, m\}$ then

$$d' = v + \frac{2\beta(1+D^{2k+1})}{(1+D)(1-D^{3k+1})}, \quad d = v + \frac{2\beta D^k(1+D^{2k+1})}{(1+D)(1-D^{3k+1})}$$

and $v_i = s_3$ equals

$$\begin{aligned} &\left(1 - \eta v - \frac{2D^{k+1}(1+D^{2k+1})}{(1+D)(1-D^{3k+1})}\right)\left(1 - \beta - v - \frac{2}{b} - \frac{2\beta D^{2k+1}(1+D^k)}{(1+D)(1-D^{3k+1})}\right) \\ &< \delta_{\infty,6} := (1 - \eta v)\left(1 - \beta - v - \frac{2}{b}\right). \end{aligned}$$

If $\{t_{2i}, t_{2i+2}\} = \{m, m\}$ we have $d, d' > v$ and

$$\begin{aligned} v_i = s_1 &\ge (1 - 2\eta + \eta v)(1 - \beta + v) \\ &= \delta_{\infty,6} + \frac{2}{b}(1 - 2\eta - \eta v) \end{aligned}$$

If $\{t_{2i}, t_{2i+2}\} = \{s, n\}$ then $w > -1/(1+D)$, $w' > 1/(1+D)$ and $d' \le v + 2\beta/(1-D)$, $d \le v - 2\beta + 2\beta D/(1-D)$ give

$$\begin{aligned} v_i = s_3 &\ge \left(1 - \eta v + \frac{2D(1-2D)}{1-D}\right)\left(1 - \beta - v - \frac{2\beta}{1+D} - \frac{4\beta D}{1-D}\right) \\ &> \delta_{\infty,6} - \frac{4\beta D}{1-D}(1 - \eta v + 2D) + \frac{2D(1-2D)}{1-D}(1 - 3\beta - v) \\ &> \delta_{\infty,6} + \frac{2D}{1-D}(1 - (5+m)\beta - D) > \delta_{\infty,6} \end{aligned}$$

for $b > 5$, giving (3.12).

For the remaining cases we have to deal with non constant t_{2i+1}.

Suppose that γ is in $S_{k,2}$, $k \geq 1$, with $m = 0$ and $b = r \geq a + 3$. If $\{t_{2i}, t_{2i+2}\} = \{n, -m\}$ then $d' = (2\beta - D)/(1 - D) - \varepsilon_{k,2}$, $d = (D - 2\beta D)/(1 - D) + D\varepsilon_{k,2}$ and $w' \leq 1/(1 - D)$, $w \leq \eta$ gives

$$\begin{aligned} v_i = s_1 &= \left(1 - 2\eta + \frac{(2D - \eta D)}{(1 - D)} - \eta\varepsilon_{k,2}\right)\left(1 - \beta + \frac{D - 2\beta D}{1 - D} + D\varepsilon_{k,2}\right) \\ &< \delta_{\infty,2} := \left(1 - 2\eta + \frac{2D - \eta D}{1 - D}\right)\left(1 - \beta + \frac{D - 2\beta D}{1 - D}\right). \end{aligned}$$

If $\{t_{2i}, t_{2i+2}\} = \{n, n\}$ then $2\beta - D \leq d, d' \leq (2\beta - D)/(1 - D)$ and

$$\begin{aligned} s_3 &\geq \left(1 - \frac{(2D - \eta D)}{1 - D}\right)\left(1 - \beta - \frac{(2\beta - D)}{1 - D}\right) \\ &= \delta_{\infty,2} + \frac{2D}{(1 - D)^2}(b - a - 3 + (7\beta - \eta - 2D - \beta D + D^2)) \end{aligned}$$

and

$$\begin{aligned} s_1 &\geq (1 - 2\eta + 2D - \eta D)(1 - \beta + 2\beta - D) \\ &> \delta_{\infty,2} - \frac{2D^2}{1 - D}(1 + \beta) + (1 - 2\eta)\frac{(2\beta - 2D)}{1 - D} > \delta_{\infty,2}, \end{aligned}$$

and (3.8) holds.

Suppose $b = a+1$ and γ is in $S_{k,4}$. Then at all points $w' \geq 1/(1-D^2)$, $w \geq D/(1-D^2)$ and $v_i = s_3$. If $t_{2i+1} = -1$ and t_{2i-1} or $t_{2i+3} = 1$ then $d' = 2\beta/(1-D^2) - D/(1-D) + \varepsilon_{k,4}$, $d = D/(1 - D) - 2\beta D/(1 - D^2) - D\varepsilon_{k,4}$ and

$$\begin{aligned} s_3 &= \left(1 - \frac{\eta D}{1 - D} + \frac{2D^2}{1 - D^2} + \eta D\varepsilon_{k,4}\right)\left(1 - \beta - \frac{2\beta}{1 - D^2} + \frac{D}{1 - D} - \varepsilon_{k,4}\right) \\ &< \delta_{\infty,4} := \left(1 - \frac{\eta D}{1 - D} + \frac{2D^2}{1 - D^2}\right)\left(1 - \beta - \frac{2\beta}{1 - D^2} + \frac{D}{1 - D}\right). \end{aligned}$$

If $t_{2i+1} = t_{2i-1} = t_{2i+3} = -1$ then $d' \leq 2\beta - D + D^2$, $d \leq -D + 2\beta D$ and

$$\begin{aligned} s_3 &\geq (1 + \eta D - 2D^2)(1 - 3\beta + D - D^2) \\ &\geq \delta_{\infty,4} + (2\eta D - 4D^2)(1 - 3\beta + D) - 2D^2(1 + D) \\ &\geq \delta_{\infty,4} + 2\eta D(1 - 6\beta + D) > \delta_{\infty,4} \end{aligned}$$

for $b \neq 4$ and (3.10) holds.

Suppose that $b = 2a+2$ and that γ is in $S_{k,7}$. Then in all cases $w' > 1/(1+D), w > -1/(1+D)$ and $v_i = s_3$. If $t_{2i+1} = -1$ and t_{2i-1} or $t_{2i+3} = 1$ then $d' = 4\beta/(1-D^2) - D/(1-D) + \varepsilon_{k,7}$ and $d = D/(1-D) - 4\beta D/(1-D^2) - D\varepsilon_{k,7}$ giving

$$\begin{aligned} s_3 &= \left(1 - \frac{\eta D}{1-D} + \frac{4D^2}{1-D^2} + \eta D\varepsilon_{k,7}\right)\left(1 - \beta + \frac{D}{1-D} - \frac{4\beta}{1-D^2} - \varepsilon_{k,7}\right) \\ &> \delta_{\infty,7} := \left(1 - \frac{\eta D}{1-D} + \frac{4D^2}{1-D^2}\right)\left(1 - \beta + \frac{D}{1-D} - \frac{4\beta}{1-D^2}\right). \end{aligned}$$

If $t_{2i+1} = t_{2i-1} = t_{2i+3} = -1$ then $d' \leq 4\beta - D + D^2$ and $d \leq -D + 4\beta D$ and

$$\begin{aligned} s_3 &> (1 + \eta D - 4D^2)(1 - 5\beta + D - D^2) \\ &> \delta_{\infty,7} - 2D^2(1+D) + (2\eta D - 8D^2)(1 - 5\beta + D) \\ &> \delta_{\infty,7} + 2\eta D(1 - 10\beta + D + 17\beta^2) > \delta_{\infty,7} \end{aligned}$$

and (3.13) holds.

Next suppose that $b = a + 2 \geq 7$ and that γ is in $S_{k,8}$. If $t_{2i+1} = -1$ and t_{2i-1} or $t_{2i+3} = -1$ then $w' \geq 1/(1+D)$, $w \geq -1/(1+D)$ and $v_i = s_3$. If $t_{2i+1} = -1$ and $\{t_{2i-1}, t_{2i+3}\} = \{-1, 1\}$ then

$$d' = \frac{\beta(3-D)}{1-D^2} - \frac{D}{1-D} + \varepsilon_{k,8}, \quad d = -\beta - \frac{D}{1-D} + 2D - \frac{\beta D(1-3D)}{1-D^2} + D^2\varepsilon_{k,8},$$

and $v_i = s_3$ equals

$$\begin{aligned} &\left(1 + D + \frac{D^2(1-3D)}{1-D^2} + \frac{\eta D^2}{1-D} - \eta D - \eta D^2\varepsilon_{k,8}\right)\left(1 - \beta - \frac{\beta(3-D)}{1-D^2} + \frac{D}{1-D} - \varepsilon_{k,8}\right) \\ &< \delta_{k,8} := \left(1 + D + \frac{D^2(1-3D)}{1-D^2} + \frac{\eta D^2}{1-D} - \eta D\right)\left(1 - \beta - \frac{\beta(3-D)}{1-D^2} + \frac{D}{1-D}\right). \end{aligned}$$

When $\{t_{2i-1}, t_{2i+3}\} = \{-1, -1\}$ we have $d' \leq 3\beta - D - \beta D + D^2$ and $d \leq -\beta - D + 3\beta D - D^2$ and

$$\begin{aligned} s_3 &\geq (1 + D + \eta D - 3D^2 + \eta D^2)(1 - 4\beta + D + \beta D - D^2) \\ &\geq \delta_{\infty,8} - 2D^2(1 + D + \eta D) + (2\eta D - 4D^2)(1 - 4\beta + D) \\ &> \delta_{\infty,8} + 2\eta D(1 - 7\beta + D) \geq \delta_{\infty,8} \end{aligned}$$

for $b \geq 7$. If $\{t_{2i-1}, t_{2i+3}\} = \{1,1\}$ then $w, w' \leq 2D$ and $d, d' \geq \beta$ give $v_i = s_1 \geq (1-2\eta+D)$ while $\delta_{\infty,8} < (1+D)(1-4\beta+D+\beta D+D^2) < 1-4\beta+2D < 1-2\eta+D$, so (3.14) holds.

Suppose now that we also allow $t_{2i+1} = -3$ with $m = 1$. For these points in S_{-5} and $S_{k,9}$ we have $d, d' \leq 3\beta - D$ and $s_3 - s_2 = 2\eta(2-d)(1-d') - 2\beta > 0$ with $s_1 - s_2 = 2(d(1-3\eta-D+\eta d) - \eta - \beta + 3D)$. Hence if $m = 1$ and γ is in S_{-5} we have $d, d' \to \frac{3\beta-3D}{1-D}$ and writing

$$s_1 - s_2 = \frac{2D}{(1-D)^2}\left(a - r - 9 + 3\beta + 12\eta - 9D + 2\beta D - \eta D\right),$$

clearly $v_i = s_1$ if $r \geq a-7$ and $v_i = s_2$ if $r \leq a-9$, giving (3.6). Suppose that γ is in $S_{k,9}$ with $m = 1$, $r = 2$ and $a \geq 9$. If $t_{2i+1} = -1$ and $\{t_{2i}, t_{2i+2}\} = \{3, -1\}$ then

$$d' = 3\beta - 3D + 3\beta D - D^2 - \frac{D^2(\beta - D)}{1-D} - \varepsilon_{k,9}D^2, \quad d = \frac{-\beta + D}{1-D} - \varepsilon_{k,9},$$

and $w' \leq 1 - \eta + D$, $w \leq -1 + \eta$ gives $v_i = s_1$ equal to

$$\left(1 - 2\eta + 3D - 3\eta D + 3D^2 - \eta D^2 - \frac{\eta D^2(\beta - D)}{1-D} - \eta D^2 \varepsilon_{k,9}\right)\left(1 - 2\beta + \frac{D(1-\beta)}{1-D} - \varepsilon_{k,9}\right)$$
$$< \delta_{k,9} := \left(1 - 2\eta + 3D - 3\eta D + 3D^2 - \eta D^2 - \frac{D^3(1-\eta)}{1-D}\right)\left(1 - 2\beta + \frac{D(1-\beta)}{1-D}\right).$$

For $t_{2i+1} = -3$ we have $d, d' \geq 3\beta - D - 2\beta D$ giving $s_1 - s_2 \geq 2D(a - 9 + 4\eta - 7\beta + d^2/\beta) > 0$ for $a \geq 9$ and $d, d' \leq 3\beta - D$ gives

$$v_i = s_2 \geq (1 - 2\eta + 3D - \eta D)(1 - 2\beta + D) > \delta_{\infty,9}.$$

Finally, when $t_{2i+1} = -1$, $t_{2i} = t_{2i+2} = 1$ we have $w, w' < \eta$ and $d, d' \geq \beta - D$ give

$$v_i = s_1 \geq (1 - 2\eta + D - \eta D)(1 - D) > 1 - 2\eta > \delta_{\infty,9}.$$

For the configurations $a = 3$, $b = 4, 5, 6$ we need also to deal with $t_{2i+1} = a$. For $t_{2i+1} = a$ we have

$$s_4 := \min\{s_2^*(2i), s_4^*(2i+1)\} = \eta(1 - \beta - d')(1 + \beta + d).$$

From the bounds $d, d' \geq -2\beta + D - \beta D - \beta D^2$ when $b = 4$, $d, d' \geq -3\beta + D - D^3$ for $b = 5$ and $d, d' \geq -2\beta$ for $b = 6$ we have $(2 - 3\beta + \beta^2) + 3d + d^2 + (d' - d)(1 + d) > 0$ and $s_1 := \min\{s_1^*(2i), s_1^*(2i+1)\} = \eta(5 - \beta + d')(1 - \beta + d) > s_4$ while $s_2^*(2i+1) = (1 + 4\eta + \eta d_{2i}^-)(1 + \beta - d_{2i+1}^+) > \eta$. Noting that switching from γ to $-\gamma$ replaces d_{2i}^- and d_{2i+1}^+ by $-d_{2i}^- - 2\beta$ and $-d_{2i+1}^+ - 2\beta$ when $t_{2i+1} = a$ (and by $-d_{2i}^-$ and $-d_{2i+1}^+$ as before otherwise) it is readily seen that we need just evaluate s_4 for the $t_{2i+1} = a$ with $t_{2i} < 0$, and $\min\{s_1, s_2, s_3\}$ for the $t_{2i+1} = -1$ (in both γ and $-\gamma$). For γ in S_{-6} and S_{-7} we $d, d' \sim -\beta$ and $d, d' \sim -\beta - \beta/(1 - D)$ and δ_{-6} and δ_{-7} are immediate. For $(a, b) = (3, 4)$ and γ in S_{-8} cutting at the $t_{2i+1} = a$ gives

$$d, d' \sim \frac{-2\beta + D - 2\beta D + aD^2}{1 - D^2}$$

with s_4 giving the value claimed for δ_{-8} while for γ in $S_{k,10}$ cutting at the $t_{2i+1} = a$ with $\{t_{2i-1}, t_{2i+3}\} = \{a, 1\}$ gives

$$d' = \left(-2\beta + aD + \frac{(-2\beta D + D^2 - 2\beta D^2 + aD^3)(1 - D^{2k})}{1 - D^2}\right)/(1 - D^{2k+1}),$$

$$d = \left(\frac{(-2\beta + D - 2\beta D + aD^2)(1 - D^{2k})}{1 - D^2} - 2\beta D^{2k} + aD^{2k+1}\right)/(1 - D^{2k+1}),$$

and the value claimed for $\delta_{k,10}$, the remaining $t_{2i+1} = a$ producing larger d and smaller d'. It is readily checked that cutting at a $t_{2i+1} = -1$ we have $\beta \leq d, d' \leq \beta(1 + D - \eta D)/(1 - D^2)$ and $\min\{s_1, s_2, s_3\} \geq \eta$ does not affect the value.

Similarly for $(a, b) = (3, 5)$ when cutting at the $t_{2i+1} = a$, $t_{2i} = t_{2i+2} = -3$, for γ in S_{-9} we have $d, d' \sim -3\beta + D - D^3(1 - 2\beta - 2\beta D + D^2)/(1 + D^4)$ while for $S_{k,11}$ the smallest d and largest d' for γ occur when t_{2i+7} or $t_{2i-5} = 1$ and $d \sim -3\beta + D - D^3(1 - 2\beta - 2\beta D + D^2)(1 - \varepsilon_{k,11})/(1 + D^4)$ and $d' \sim -3\beta + D + D^3(1 - 2\beta - 2\beta D + D^2)(1 + D^4\varepsilon_{k,11})/(1 + D^4)$. These give the values claimed for δ_{-9} and $\delta_{k,11}$. When $t_{2i+1} = a$ and $t_{2i} = t_{2i+2} = -1$ we have $d' \leq -\beta + D - 2\beta D - 2\beta D^2 + D^3 + D^4$, $d \geq -\beta - D + 2\beta D + 2\beta D^2 - D^3 - D^4$ and $s_4 > \eta\left(1 - \left(\frac{2\beta - D - D^3 + 2\beta D^3}{1 - D^4}\right)^2\right)$ (the value claimed for $\delta_{0,11}$ and the largest of those claimed). For the $t_{2i+1} = -1$ we have $-D^2 \leq d \leq D^2$ and $2\beta + \beta D \leq d' \leq 2\beta + 2\beta D$ and $\min\{s_1, s_2, s_3\}$ is also larger. When $k \geq 1$ the $t_{2i+1} = a$, $t_{2i} = t_{2i+2} = -3$ in $-\gamma$ have $d \geq$

$-3\beta+D-D^3+2\beta D^3+2\beta D^4-D^5-D^7$ and $d' \le -3\beta+D-D^3+2\beta D^3+2\beta D^4-D^5+D^7$ and s_4 is certainly greater than the value claimed for $\delta_{1,11}$.

For $(a,b)=(3,6)$ and γ in $S_{k,12}$ cutting at the $t_{2i+1}=a$ gives

$$d, d' \sim -\beta - \frac{\beta(1-\eta+D)(1-D^{k+1})}{(1-D)(1-D^{k+2})}$$

and the value claimed for $\delta_{k,12}$, while the remaining $t_{2i+1}=-1$ have $\beta \le d, d' \le 2\beta$ and $\min\{s_1,s_2,s_3\} \ge \eta$. ■

Proof of Lemma 3 ($a = 2$)

We assume that the $a_{2i-1}=a=2$ and the $a_{2i}=b\ge 5$. Using the forwards-backwards symmetry in the expansions of the γ in question it is straightforward to see that it is enough to check

$$\begin{aligned} s_1^*(2i-1) &= (1-\eta+d^-_{2i-1})(1-\beta+d^+_{2i-1}),\\ s_2^*(2i-1) &= (1+\eta+d^-_{2i-1})(1+\beta-d^+_{2i-1}),\\ s_4^*(2i-1) &= (1+\eta-d^-_{2i-1})(1+\beta+d^+_{2i-1}), \end{aligned}$$

and the corresponding functions, $s_1^*(2i-1)', s_2^*(2i-1)', s_4^*(2i-1)'$ say, for $-\gamma$.

Now if $t_{2i-1}=0$ then going from γ to $1-\alpha-\gamma$ gives $(d^-_{2i-1}, d^+_{2i-1})' = (-d^-_{2i-1}, -d^+_{2i-1})$, and $s_1^*(2i-1)', s_2^*(2i-1)', s_4^*(2i-1)'$ merely add the function

$$s_3^*(2i-1) = (1-\eta-d^-_{2i-1})(1-\beta-d^+_{2i-1}).$$

Writing $s_1^*(2i-1), s_3^*(2i-1) = \eta(1-\beta\pm d^-_{2i-2})(1-\beta\pm d^+_{2i-1})$ clearly

$$v_i := \min\{s_1^*(2i-1), s_3^*(2i-1)\} = \eta(1-\beta-v')(1-\beta+v)$$

where

$$v' := \max\{-d^-_{2i-2}, d^+_{2i-1}\}, \qquad v := \min\{-d^-_{2i-2}, d^+_{2i-1}\}.$$

Now for the γ in question we have $|d^+_{2i-1}| \le \beta, |d^-_{2i-1}| \le D$, giving $s_2^*(2i-1), s_4^*(2i-1) > 1$ (certainly larger than δ_k), so it is enough just to evaluate v_i when $t_{2i-1}=0$.

Now if $t_{2i-1} = a$ then

$$s_4^*(2i-1) = \eta(1-\beta-d_{2i-2}^-)(1+\beta+d_{2i-1}^+)$$

while going from γ to $1-\alpha-\gamma$ gives $(d_{2i-2}^-, d_{2i-1}^+)' = (-2\beta - d_{2i-2}^-, -2\beta - d_{2i-1}^+)$,

$$s_4^*(2i-1)' = \eta(1+\beta+d_{2i-2}^-)(1-\beta-d_{2i-1}^+),$$

and

$$u_i := \min\{s_4^*(2i-1), s_4^*(2i-1)'\} = \eta(1+\beta+d)(1-\beta-d'),$$

where

$$d' := \max\{d_{2i-2}^-, d_{2i-1}^+\}, \qquad d := \min\{d_{2i-2}^-, d_{2i-1}^+\}.$$

Suppose that the t_i in the expansion of γ have period $a, -t$. Then $d_{2i-1}^- \sim (a\eta - tD)/(1-D)$, $d_{2i-2}^-, d_{2i-1}^+ \sim (-t\beta + aD)/(1-D)$ and $d_{2i-1}^+(1+\eta) \le d_{2i-1}^-(1+\beta)$ giving $s_4^*(2i-1) \le s_2^*(2i-1)$. Now if $t \ge 2$ then it is easily seen that ${d_{2i-1}^+}', {d_{2i-1}^-}' \ge d_{2i-1}^+, d_{2i-1}^-$ so that

$$\min\{s_1^*(2i-1), s_1^*(2i-1)'\} = s_1^*(2i-1) \sim \eta\left(\left(2 - 2\beta - \frac{(t-2)\beta}{1-D}\right)^2 - 1\right)$$

while $u_i \sim \eta\left(1 - \left(\frac{(t-2)\beta}{1-D}\right)^2\right)$ and $\delta_{0,t}$ will be the minimum of these two expressions. Note that the second expression is smaller than the first when $t \le b - \sqrt{2b-4}$ and greater than δ_∞ when $(t-2) \le \sqrt{2b-4}$ (for the finite number of b and t with $(t-2) \le \sqrt{2b-4}$ but $t > b - \sqrt{2b-4}$ one can check numerically that $\delta_{0,t} \ge \delta_\infty$ except for $(b,t) = (6,4)$ or $(7,5)$).

For the remaining γ if $t_{2i-1} = a$ we have $d_{2i-1}^+ < aD$, $d_{2i-1}^- \ge 1-\eta$ so $s_4^*(2i-1) < s_2^*(2i-1)$. Also the rough bounds $d_{2i-2}^-, d_{2i-1}^+ \ge (-4\beta + aD)/(1-D)$ when $b \ge 8$ and $d_{2i-2}^-, d_{2i-1}^+ \ge (-3\beta + aD)/(1-D)$ when $b = 7$, and $d_{2i-2}^-, d_{2i-1}^+ \ge (-4\beta + aD - 2\beta D + aD^2)/(1-D^2)$, $(-2\beta + aD - 4\beta D + aD^2)/(1-D^2)$ for γ in S_{2k} and $d_{2i-2}^-, d_{2i-1}^+ \ge -2\beta$ for γ in S_{2k+1} when $b = 6$, give $\beta < \eta(1+d_{2i-2}^-)(1+d_{2i-1}^+) = (1+d_{2i-1}^+)(d_{2i-1}^- - \eta)$ and hence $s_4^*(2i-1) \le s_1^*(2i-1)$ when $b \ge 6$. When $b = 5$ we have $d_{2i-1}^- \ge (a\eta - 3D)/(1-D)$, $d_{2i-1}^+ \ge (-3\beta + aD)/(1-D)$ and $s_1^*(2i-1) \ge 2\eta/3$ (larger than the values claimed for

the δ_k) and so can be similarly ignored. Hence for the $t_{2i-1} = a$ we need only consider u_i.

For period $(a,-4)(a,-2)^k$ the smallest d and largest d' occur simultaneously when $t_{2i} = -4$ (or $t_{2i-2} = -4$) and

$$d' = d^-_{2i-2} \sim \frac{-2\beta + aD}{1-D} - \frac{2\beta D^k}{1-D^{k+1}} = -\beta - \frac{2\beta D^k}{1-D^{k+1}}$$
$$d = d^+_{2i-1} = \frac{-2\beta + aD}{1-D} - \frac{2\beta}{1-D^{k+1}} = -3\beta - \frac{2\beta D^{k+1}}{1-D^{k+1}}$$

with u_i asymptotically giving the value claimed for δ_{2k}, b even.

Similarly for period $(a,-3)(a,-3,a,-1)^k$ the smallest d and largest d' again occur simultaneously when $..., t_{2i-2}, t_{2i-1} = ... - 3, a, -3, a$ and

$$\begin{aligned} d = d^-_{2i-2} \sim & \frac{aD}{1-D} + \left(-3\beta + (-3\beta D - \beta D^2)\frac{(1-D^{2k})}{1-D^2} \right) /(1-D^{2k+1}) \\ & = -2\beta - \frac{1}{2}\beta^2 - \frac{2\beta D^{2k+2}}{(1+D)(1-D^{2k+1})} \\ d' = d^+_{2i-1} = & \frac{aD}{1-D} + \left((-\beta - 3\beta D)\frac{(1-D^{2k})}{1-D^2} - 3\beta D^{2k} \right) /(1-D^{2k+1}) \\ & = -\frac{1}{2}\beta^2 - \frac{2\beta D^{2k}}{(1+D)(1-D^{2k+1})} \end{aligned}$$

with this u_i asymptotically giving the value claimed for δ_{2k}, b odd.

For period $(a,-3,a,-1)$ we clearly have $d' \sim (-\beta + aD - 3\beta D + aD^2)/(1-D^2)$ and $d \sim (-3\beta + aD - \beta D + aD^2)/(1-D^2)$, giving δ_{-2}. Likewise for γ in S_{-1}, b odd, the largest and smallest d', d occur when cutting the sequence $..., t_{2i-1}|t_{2i}, ...$ at the point $-3, a, -3, a, -3, a| - 1, a$ and

$$d' = d^+_{2i-1} = \frac{-\beta - 3\beta D - 3\beta D^2 - 3\beta D^3}{1-D^4} + \frac{aD}{1-D} = -\frac{\beta D(1 + D + D^2 - D^3)}{1-D^4}$$
$$d = d^-_{2i-2} \sim \frac{-3\beta - 3\beta D - 3\beta D^2 - \beta D^3}{1-D^4} + \frac{aD}{1-D} = -2\beta - \frac{\beta D(1 + D - D^2 + D^3)}{1-D^4},$$

with u_i giving the stated value of δ_{-1}.

Now if γ has period $(a,0,0,-2)(a,-2)^k$ then cutting at a $t_{2i-1} = a$ gives

$$d' \le \frac{-2\beta D + aD^2}{1-D^2} = -\frac{1}{2}\beta^2,$$
$$d \ge \frac{-2\beta + aD^2}{1-D^2} = -2\beta + \frac{1}{2}\beta^2,$$

and the $u_i \geq \eta(1-\beta+\frac{1}{2}\beta^2)^2$, greater than the value claimed for δ_{2k+1}, b even.

Now cutting at the $t_{2i-1}=0$ gives

$$d^-_{2i-2}\sim\frac{aD-2\beta D}{1-D}+\frac{(-aD^2+2\beta D^2)D^k}{1-D^{k+2}}=\frac{\beta(1-D^{k+1})}{1-D^{k+2}}$$
$$d^+_{2i-1}=\frac{-2\beta+aD}{1-D}+\frac{(2\beta D-aD^2)D^k}{1-D^{k+2}}=-\frac{\beta(1-D^{k+1})}{1-D^{k+2}}$$

and v_i gives the valued claimed for δ_{2k+1}, b even.

Finally for period $(a,-1,0,-1)(a,-1,a,-3)^k$ cutting at a $t_{2i-1}=a$ we have

$$d\geq\frac{-3\beta+aD-\beta D-\beta D^2+aD^3-\beta D^3+aD^4}{1-D^4}=-2\beta+\frac{\beta(1-D)D^3}{1-D^4}$$
$$d'\leq\frac{-\beta+aD-\beta D-\beta D^2+aD^3-3\beta D^3+aD^4}{1-D^4}=-\frac{\beta(1-D)D^3}{1-D^4}$$

and $u_i\geq\eta(1-\beta)^2$ is larger than the value claimed for δ_{2k+1}. Lastly cutting at the $t_{2i-1}=0$ we have

$$d^+_{2i-1}=-\beta+\frac{aD-\beta D+aD^2-3\beta D^2}{1-D^2}-\frac{(aD-2\beta D)D^{2k+1}}{1-D^{2k+2}}=\frac{1}{2}\frac{\beta^2(1-D^{2k})}{1-D^{2k+2}}$$
$$d^-_{2i-2}\sim\frac{-\beta+aD-3\beta D+aD^2}{1-D^2}+\frac{(2\beta-aD)D^{2k+1}}{1-D^{2k+2}}=-\frac{1}{2}\frac{\beta^2(1-D^{2k})}{1-D^{2k+2}}$$

and v_i asymptotically gives the value claimed for δ_{2k+1}, b odd. ■

4. Proof of Theorem 1 (even $a\geq 4$)

Writing

$$u=\begin{cases}1-\eta-2\beta+D, & \text{if } b \text{ is odd},\\ 1-\eta-3\beta+3D, & \text{if } b \text{ is even},\end{cases}$$

we note that $\delta_\infty\geq(1-2\beta)(1-\eta-D)>u$ when b is odd and $\delta_{\infty,1}\geq(1-\eta)(1-3\beta)=u$ when b is even.

Since $u>\eta$ we can by (1.2) rule out $t_i=a_i$ infinitely often. In particular we can assume that going from γ to $1-\gamma$ merely replaces the t_i with $-t_i$. Since $1-4\eta+D<u$ we can further assume from (1.3) that $|t_{2i-1}|=0,2$ for all i, and since $1-2\eta+D<u$ when b is odd and $\eta>2\beta$ or when b is even and $\eta\geq 3\beta-2D$, that $t_{2i-1}=0$ when $b\geq 2a+1$ is odd or $b\geq 3a-2$ is even. Likewise we can assume that $|t_{2i}|=1,3$ when

$b \le 2a-1$ is odd and $|t_{2i}| = 0, 2$ or 4 when $b \le 3a+2$ is even and $|t_{2i}| = 0, 2$ when $b = a+2$ (else $M^*(\alpha,\gamma)$ is less than $1-5\beta+D < u$ or $1-6\beta+D < u$ or $1-4\beta+D < u$ respectively). From the bound

$$\delta_{\infty,4} \ge (1-\eta+2D-2\eta D)(1-3\beta+2D-2\beta D) \ge 1-3\beta-\eta+7D-4\eta D-8\beta D$$

and numerically for $\delta_{\infty,6}$ we similarly obtain $|t_{2i}| = 0, 2$ for $b = a+4$, $a \ge 8$.

Now if $t_{2i+1} = 0$ and $d := \max\{|d^+_{2i+1}|, |d^-_{2i}|\} \ge 2\beta$ when b is odd or $d \ge 3\beta$ when b is even then

$$\min\{s_1^*(2i+1), s_3^*(2i+1), s_1^*(2i), s_3^*(2i)\} \le (1-\beta-d)(1-\eta+\eta d) \tag{4.1}$$

is less than u. Hence we can assume if $t_{2i+1} = 0$ then $t_{2i}, t_{2i+2} = \pm 1$ or $0, \pm 2$ as b is odd or even.

If $t_{2i+1} = -2$ then (with at most a finite number of exceptions) $t_{2i}, t_{2i-2} \ge 0$ else (since the $t_{2i} \le 4$) the $s_1^*(2i)$ or $s_1^*(2i+1)$ give

$$\begin{aligned} M^*(\alpha,\gamma) &\le \left(1-3\eta+\frac{4D+2\eta D}{1-D}\right)\left(1-3\beta+\frac{2D+4\beta D}{1-D}\right) \\ &\le 1-3\eta-3\beta+15D-4\eta D < u \end{aligned}$$

for b even and

$$\begin{aligned} M^*(\alpha,\gamma) &\le \left(1-3\eta+\frac{3D+2\eta D}{1-D}\right)\left(1-2\beta+\frac{2D+3\beta D}{1-D}\right) \\ &\le 1-3\eta-2\beta+11D-4\eta D < u \end{aligned}$$

for b odd (in particular if $t_{2i} = 3$ or 4 we can assume that $t_{2i\pm1} = -2$).

Suppose first that b is odd. If $t_{2i+1} = -2$ and $t_{2i} \ne 3$ then $t_{2i} = 1$ and

$$s_1^*(2i) \le \left(1-3\eta+\frac{3D-2\eta D}{1-D}\right)\left(1-\beta+\frac{\beta}{1-D}\right) < 1-3\eta+3D-2\eta D+2\beta D \le u.$$

Hence for b odd we can assume that the expansion of γ either has period C_3 or C_3' or consists solely of blocks A_1', A_1. We suppose the expansion consists only of blocks A_1, A_1' and assume that γ is not in S_0 so that γ (or its negative) has infinitely many

blocks $t_{2i-1},...,t_{2i+3}=0,-1,0,-1,0$. We can rule out blocks $A_1A_1A_1$ or $A_1'A_1'A_1'$ since $t_{2i+4}=-1$ gives

$$s_1^*(2i+1)\leq\left(1-\eta-D+\frac{D^2}{1-D}\right)\left(1-2\beta-\beta D+\frac{\beta D^2}{1-D}\right)<(1-2\beta)(1-\eta-D),$$

less than δ_∞, and we can rule out $t_{2i+4},t_{2i+5},...$ (or $t_{2i-2},t_{2i-3},...$) of the forms $1,0,-1,0,-1,...$ or $1,0,-1,0,1,0,-1,...$ else

$$d_{2i+1}^+\leq-\beta+\beta D-\beta D^2+\frac{\beta D^3(1-D+D^2)}{1-D^3}=-\frac{\beta(1-D)(1+2D^2)}{1+D^2}-\frac{2\beta D^4(1-D-D^3)}{(1+D^2)(1-D^3)},$$

$$d_{2i+1}^-\leq\frac{-D+D^2+D^3}{1-D^3}=-\frac{D(1-D)}{1+D^2}+\frac{2D^5}{(1+D^2)(1-D^3)},$$

and $s_1^*(2i+1)\leq\delta_\infty-\frac{2\beta D^4}{(1+D^2)(1-D^3)}((1-D-D^3)(1-\eta-D)-\eta)<\delta_\infty$. Hence either γ has period $A_1A_1A_1'A_1'$ or γ (or its negative) has infinitely many blocks $t_{2i-1},...,t_{2i+11}=0,-1,0,-1,0,1,0,-1,0,1,0,1,0$ and writing $d_{2i+1}^-=-D(1-D)/(1+D^2)+w^-$ and $d_{2i+8}^+=D(1-D)/(1+D^2)-w^+$ we have

$$s_1^*(2i+1)=\left(1-\eta-\frac{D(1-D)}{1+D^2}+w^-\right)\left(1-2\beta+\beta D-\frac{\beta D^2(1-D)}{1+D^2}-\beta D^3w^+\right)$$
$$s_3^*(2i+8)=\left(1-\eta-\frac{D(1-D)}{1+D^2}+w^+\right)\left(1-2\beta+\beta D-\frac{\beta D^2(1-D)}{1+D^2}-\beta D^3w^-\right).$$

Now $w=\min\{w^+,w^-\}\geq0$ else the minimum of these is certainly less than

$$\left(1-\eta-\frac{D(1-D)}{1+D^2}+w\right)\left(1-2\beta+\beta D-\frac{\beta D^2(1-D)}{1+D^2}-\beta D^3w\right)<\delta_\infty.$$

Hence we can assume that the sequence for γ consists of blocks $A_1A_1'(A_1A_1A_1'A_1')^{l_i}$. Now if our block occurs inside a block

$$...0,1,0,-1(0,1,0,1,0,-1,0,-1)^l0,1,0,-1,0(1,0,1,0,-1,0,-1,0)^m1,0,-1,0...$$

with $l>m$ then $w^-\leq2D^{1+4l}\leq2D^{4m+5}$ and $-w^+\leq-2D^{4m+1}(1-D-D^2)$ and $s_1^*(2i+1)\leq\delta_\infty-2\beta D^{4m+4}((1-\eta-D)(1-D-D^2)-\eta)<\delta_\infty$. Hence we need only worry about the equal lengthed blocks of S_k. So the only values above δ_∞ come from

these δ_k, $k \geq 0$, and δ_{-1}, plus δ_{-2} when $b < 2a$ and $\delta_{-2} > \delta_\infty$ (one tediously checks that this and $\delta_{-2} > \delta_0$ occur as stated).

Suppose now that b is even. From the above we need only consider blocks A_0, A_2, C_0, C_2, C_4 or their negatives.

We also note that if $t_{2i+1} = -2$ and $t_{2i} = 0$ (or $t_{2i+2} = 0$) then

$$\begin{aligned} s_1^*(2i) &\leq \left(1 - \beta + \frac{2D}{1-D}\right)\left(1 - 3\eta + \frac{4D - 2\eta D}{1-D}\right) \\ &\leq 1 - 3\eta - \beta + 9D = u - 2D(b - a - 3 - \eta + \beta), \end{aligned}$$

is less than u for $b \geq a + 4$ and

$$s_1^*(2i) \leq \left(1 - \beta + \frac{2D}{1-D}\right)\left(1 - 3\eta + \frac{2D}{1-D}\right) \leq 1 - 3\eta - \beta + 7D - 6\eta D,$$

for $b = a + 2$ enabling us to rule out blocks C_0.

If $t_{2i-1} = -2$ and $t_{2i} \leq 2$ then

$$\begin{aligned} s_1^*(2i-1) &\leq \left(1 + \beta + \frac{2\beta D}{1-D}\right)\left(1 - 3\eta + \frac{4D - 2\eta D}{1-D}\right) \\ &\leq 1 - 3\eta + \beta + D + 6\beta D - 2\eta D = u - 2D(b - 2a + 1 - \beta). \end{aligned}$$

Hence for $b \geq 2a$ we can assume that either the expansion of γ eventually has period C_4 or that $t_{2i-1} = 0$ for all i.

Suppose first that $b \geq 2a$. Now infinitely many blocks of the form ${A'}_2{A'}_2$ or ${A'}_2A_0{A'}_2$ or ${A'}_2A_0A_0$ (or their negatives or reverses) would lead to

$$M^*(\alpha, \gamma) \leq \left(1 - 3\beta + \frac{2\beta D}{1-D}\right)\left(1 - \eta + \frac{2D^3}{1-D}\right) \leq \delta_{\infty,1}.$$

Hence if γ does not have period C_4 or A_0 or $A_2{A'}_2$ then we can assume that its expansion consists of infinitely many A_0 separated by alternating A_2 and ${A'}_2$. We suppose that $t_{2i} = 0$, $t_{2i-2} = -2$, $t_{2i+2} = 2$ and write $d^+_{2i+1} = \frac{2\beta}{1+D} + w^+$, $d^-_{2i-2} = -\frac{2\beta}{1+D} - w^-$, so that

$$\begin{aligned} s_3^*(2i+1) &= \left(1 - \beta - \frac{2\beta}{1+D} - w^+\right)\left(1 - \eta + \frac{2D^2}{1+D} + \eta D w^-\right) \\ s_1^*(2i-2) &= \left(1 - \beta - \frac{2\beta}{1+D} - w^-\right)\left(1 - \eta + \frac{2D^2}{1+D} + \eta D w^+\right) \end{aligned}$$

We can clearly assume that $w = \max\{w^+, w^-\} < 0$ else the minimum of these will be at most

$$\left(1 - \beta - \frac{2\beta}{1+D} - w\right)\left(1 - \eta + \frac{2D^2}{1+D} + \eta D w\right) \leq \delta_{1,\infty} - w(1 - \eta - \eta D) < \delta_{\infty,1}.$$

Hence the $t_{2i} = 0$ occur in the middle of blocks

$$A_0(A_2A'_2)^m A_0(A_2A'_2)^l A_0.$$

Now if $l > m$ then $w^- \leq -2\beta(1-2D)D^{2m}$ and $w^+ \geq -2\beta D^{2l} \geq -2\beta D^{2m+2}$ giving

$$s_3^*(2i+1) \leq \delta_{\infty,1} - 2D^{2m+2}((1-2D)(1-3\beta) - \beta) < \delta_{\infty,1}.$$

Hence this leaves only the periodic case $A_0(A_2A'_2)^l$ and the result follows (after checking that $\delta_{0,4} > \delta_{1,\infty}$ iff $2a \leq b \leq 3a - 6$).

Suppose next that $b = 2a - 2$, $a \geq 8$ and note the bound

$$\delta_{\infty,2} \geq 1 - 3\eta + \beta + D - 2\eta D + 2\beta D.$$

If $t_{2i-1} = -2$, $t_{2i} = 2$ and either $t_{2i+1} = -2$ or $t_{2i+1}, t_{2i+2}, t_{2i+3} = 0, -2, 0$ then

$$s_1^*(2i-1) \leq \left(1 - 3\eta + \frac{4D - 2\eta D}{1-D}\right)\left(1 + \beta - 2\beta D + \frac{2\beta D^2}{1-D}\right) < \delta_{\infty,2},$$

and if we have blocks $t_{2i-2}, t_{2i-1}, t_{2i} = 2, -2, 2$ then

$$s_1^*(2i-1) \leq \left(1 - 3\eta + \frac{2D}{1-D}\right)\left(1 + \beta + \frac{2\beta D}{1-D}\right) \leq 1 - 3\eta + \beta - D + 4\beta D < \delta_{\infty,2},$$

and if we have blocks $t_{2i-2}, ..., t_{2i+1} = 4, -2, 2, 0$, with $t_{2i+2} \geq 0$ then

$$\begin{aligned} s_3^*(2i) &\leq \left(1 - 3\beta + 2D - 4\beta D + \frac{2D^2}{1-D}\right)\left(1 - \eta + \frac{2\eta D}{1-D}\right) \\ &\leq 1 - 3\beta - \eta + 5D - 4\beta D + 4\eta D^2 < \delta_{\infty,2}. \end{aligned}$$

Hence if we have infinitely many $t_{2i-1} = -2$ then either γ has period C_4 or we have infinitely many blocks $t_{2i-2}, ..., t_{2i+4} = 4, -2, 2, 0, -2, 2, -4$. Hence we have

$$\begin{aligned} s_1^*(2i-1) &= (1 - 3\eta + \eta w^-)(1 + \beta - 2\beta D + 2D^2 - D^2 w^+) \\ s_3^*(2i+2) &= (1 - 3\eta + \eta w^+)(1 + \beta - 2\beta D + 2D^2 - D^2 w^-) \end{aligned}$$

where $w^+, w^- \leq (4\beta - 2D)/(1-D)$ and $\min\{s_1^*(2i-1), s_3^*(2i+2)\} \leq \delta_{\infty,2}$. Hence it remains only to deal with cases where $t_{2i-1} = 0$ for all i as for $b \geq 2a$. Since $\delta_{0,5} \leq 1 - \eta - 3\beta + 5D - 4\beta D < \delta_{\infty,2}$ for $b = 2a-2$, these only add $\delta_{0,1}$.

Suppose next that $b = 2a-4$, $a \geq 10$. We note the upper bound

$$\delta_{\infty,3} \geq (1 - \eta - 2\eta D + 2D)(1 - 3\beta + 2D - 4\beta D) \geq 1 - 3\beta - \eta + 7D - 4\eta D - 10\beta D,$$

and rule out the configuration $t_{2i-2}, t_{2i-1} = -2, 0$ with $t_{2i} \leq 0$, since this leads to

$$s_1^*(2i-2) \leq (1 - 3\beta + 2D)\left(1 - \eta + \frac{2D^2}{1-D}\right) \leq 1 - \eta - 3\beta + 5D < \delta_{\infty,3},$$

and $t_{2i-2}, ..., t_{2i+1} = 2, -2, 2, -2$, since for $a \geq 10$ this leads to

$$\begin{aligned} s_1^*(2i-1) &\leq (1 - 3\eta + 2D)\left(1 + \beta - 2D + \frac{4\beta D - 2D^2}{1-D}\right) \\ &\leq 1 + \beta - 3\eta - 3D + 6\eta D + 6\beta D < \delta_{\infty,3}. \end{aligned}$$

Suppose that we have infinitely many blocks $t_{2i-2}, t_{2i-1}, t_{2i} = -2, 0, 2$. If $t_{2i-3} = 0$, or $t_{2i-3}, t_{2i-4} = 2, -4$ then (4.1) with $d \geq 2\beta - \frac{(2D - 4\beta D + 2D^2 - 2\beta D)}{1-D^2}$ (since we have excluded blocks C_2C_2 or C_0) gives $s_1^*(2i-2)$ or $s_3^*(2i-1)$ less than $\delta_{\infty,3}$. So we need only include periods A_0 or $A_2C_2A_2'C_2'$ or γ consisting solely of blocks C_2 and C_4 with no blocks C_2C_2. Hence either γ has period C_4 or C_2C_4 or γ consists of blocks $..., b_{2i}|b_{2i+1}, ...$ of the form $..., C_4(C_4C_2)^k|C_4C_4(C_2C_4)^lC_4, ...$ Hence, writing $d_{2i+1}^- = \frac{-2\eta}{1-D} + \frac{2D+4D^2}{1-D^2} + w^-$ and $d_{2i+4}^+ = \frac{-2\eta}{1-D} + \frac{2D+4D^2}{1-D^2} + w^+$, if $0 \leq l < k$ we have $w^- \leq \frac{2D^{2k+1}}{1-D^2} \leq \frac{2D^{2l+3}}{1-D^2}$ and $w^+ \geq \frac{2D^{2l+1}}{1-D^2}(1-D)$ and $s_2^*(2i+1)$ is

$$\begin{aligned} &\left(1 - \eta - \frac{2\eta D}{1-D} + \frac{2D+4D^2}{1-D^2} + w^-\right)\left(1 - 3\beta + \frac{2D}{1-D} - \frac{(4\beta D + 2\beta D^2)}{1-D^2} - \beta D w^+\right) \\ &\leq \delta_{\infty,3} - \frac{2\beta D^{2l+2}}{1-D^2}((1-\eta)(1-D) - \eta) < \delta_{\infty,3}, \end{aligned}$$

leaving only the periodic elements of $S_{k,3}$.

Suppose now that $b \leq 2a-6$ or $a = 6, b = 8$. If $t_{2i+1} = 0$ and t_{2i} or $t_{2i+2} = \pm 2$ infinitely often then (4.1) with $d \geq (2\beta - 2D)/(1-D)$ gives $M^*(\alpha, \gamma) \leq \delta_{\infty,4}$.

Hence when $b \le a+4 \le 2a-6$, or $a=6, b=8$, the only values above $\delta_{\infty,4}$ come from periods A_0 or C_2. When $a+6 \le b \le 2a-6$ we need also to consider γ consisting of C_2 and C_4 with infinitely many C_4. Suppose we have blocks $...,b_{2i-2}|b_{2i-1},...$ of the form $...C_4C_2^l|C_4C_2^mC_4...$ with $l>m\ge 0$ then, writing $d^-_{2i-1}=\frac{-2\eta+2D}{1-D}+w^-$ and $d^+_{2i-1}=4\beta-\frac{2D(1-\beta)}{1-D}+\beta w^+$, the bounds $w^-\le 2D^{l+1}/(1-D)\le 2D^{m+2}/(1-D)$ and $w^+\ge 2D^{m+1}$ give

$$\begin{aligned} s_2^*(2i-1) &= \left(1-\eta+\frac{2D(1-\eta)}{1-D}+w^-\right)\left(1-3\beta+\frac{2D(1-\beta)}{1-D}-\beta w^+\right)\\ &\le \delta_{\infty,4}-2\beta D^{m+1}((1-\eta)-\eta)\le \delta_{\infty,4}. \end{aligned}$$

Hence it is enough to consider the periodic elements in $\delta_{k,4}$, $k\ge 0$.

It remains to deal with the left-over cases $(a,b)=(4,6)$, $(6,10)$ and $(8,12)$.

When $(a,b)=(4,6)$ we can rule out blocks $t_{2i-3},...,t_{2i}=0,-2,0,-2$ else

$$s_1^*(2i-2)\le\left(1-3\beta+\frac{2\beta D}{1-D}\right)\left(1-\eta-2D+\frac{2\eta D-2D^2}{1-D}\right)<\delta_{\infty,1}$$

and $t_{2i-2},...,t_{2i+3}=-2,0,0,0,-2,0$, else

$$s_1^*(2i+1)\le\left(1-3\beta+\frac{2\beta D}{1-D}\right)\left(1-\eta-2D^2+2\eta D^2\right)<\delta_{\infty,1}.$$

Hence if we have blocks $t_{2i-1},t_{2i}=-2,2$, with $t_{2i+1}=-2$, or with $t_{2i+1}=0$ and $t_{2i+2}\le 0$, then

$$s_1^*(2i-1)\le\left(1-3\eta+\frac{2D}{1-D}-\frac{2\eta D^2}{1-D^2}\right)\left(1+\beta+\frac{2\beta D^2}{1-D}-\frac{2D^3}{1-D^2}\right)<\delta_{\infty,1}.$$

So either γ has period A_2C_2 or we can assume that $t_{2i-1}=0$ for all i (in which case the proof follows exactly as for $b\ge 2a$).

For $(a,b)=(6,10)$ observe that we can have no blocks $t_{2i},t_{2i+1},t_{2i+2}=-4,2,-4$ else

$$s_1^*(2i-1)\le\left(1-5\beta+\frac{2D-2\beta D}{1-D}\right)\left(1+\eta-4D+\frac{2\eta D-2D^2}{1-D}\right)<\delta_{\infty,7},$$

and if we have $t_{2i-1}=-2$ then t_{2i-2} or $t_{2i}=4$ else

$$s_1^*(2i-1)\le\left(1-3\eta+\frac{2D}{1-D}\right)\left(1+\beta+\frac{2\beta D}{1-D}\right)<\delta_{\infty,7}.$$

Likewise we can rule out blocks $t_{2i-1},...,t_{2i+1}=0,-2,0$ and $t_{2i+2}\leq 0$ else

$$s_1^*(2i)\leq\left(1-3\beta+\frac{2\beta D}{1-D}\right)\left(1-\eta+\frac{2D^2}{1-D}\right)<\delta_{\infty,7}.$$

Hence if γ has infinitely many $t_{2i-1}=\pm 2$ but not period C_2C_4 then we must have infinitely many blocks $t_{2i-2},...,t_{2i+1}=-4,2,-2,0$ and $s_1^*(2i-2)\leq\delta_{\infty,7}$. If $t_{2i-1}=0$ for almost all i then all but period A_2A_2' and A_0 have been ruled out.

For $(a,b)=(8,12)$ we first rule out $t_{2i-1},...,t_{2i+3}=-2,2,-2,2,-2$ else

$$s_1^*(2i+1)\leq\left(1-3\eta+2D-2\eta D+\frac{2D^2}{1-D}\right)\left(1+\beta-2D+\frac{2\beta D}{1-D}\right)<\delta_{\infty,6}.$$

If $t_{2i-1},t_{2i}=0,-2$ then $t_{2i+1},t_{2i+2}=2,-2$ else

$$s_1^*(2i-1)\leq\left(1-\eta+\frac{2D}{1-D}\right)\left(1-3\beta+\frac{2\beta D}{1-D}\right)<\delta_{\infty,6}$$

and $t_{2i-2},t_{2i-3}=2,-2$ else

$$s_1^*(2i-1)\leq\left(1-\eta+\frac{2D^2}{1-D}\right)\left(1-3\beta+\frac{2D-2\beta D}{1-D}\right)<\delta_{\infty,6}.$$

Likewise if $t_{2i-1},t_{2i-2}=0,-2$ then $t_{2i-4},...,t_{2i+1}=-2,2,-2,0,2,-2$. Hence blocks A_2' occur inside blocks $C_2A_2'C_2'$ and either γ has period $A_2C_2A_2'C_2'$ or we have infinitely many blocks $b_{2i-1},...,b_{2i+4}=C_2'A_2C_2$ with $b_{2i+5}b_{2i+6}=C_2$ and, writing

$$d_{2i+3}^+,-d_{2i-2}^-=\frac{2\beta+2\beta D-2D}{1+D^2}+w^+,w^-$$

we have

$$s_1^*(2i+3)=\left(1+\beta-2D\frac{(1-\beta+\beta D)}{1+D^2}+w^+\right)\left(1-3\eta+2D-2D^2+2\eta D^2-2D^3\frac{(1-\eta+D)}{1+D^2}-\eta D^2w^-\right),$$

with a similar expression for $s_3^*(2i-2)$ with w^+ and w^- interchanged. Clearly then $w=\min\{w^+,w^-\}>0$ else the minimum of $s_1^*(2i+3)$ and $s_3^*(2i-2)$ will be less than $\delta_{\infty,6}+w((1-3\eta)-\eta D^2(1+\beta))<\delta_{\infty,6}$. Now if this block $C_2'A_2C_2$ occurs inside a block

$$...C_2C_2(A_2'C_2'A_2C_2)^lA_2'C_2'C_2'A_2C_2C_2(A_2'C_2'A_2C_2)^kA_2'C_2'C_2'...$$

with $k > l \geq 0$ or

$$...A_2'C_2'A_2C_2C_2(A_2'C_2'A_2C_2)^kA_2'C_2'C_2'...$$

with $k \geq 0$, then $w^+ \leq 2D^{4+4k}$ and $w^- \geq 2D^{4+4l}(1-2\beta)$ or $w^- \geq 2D(1-2\beta)$ respectively, and $w^- \geq 2D^{1+4k}(1-2\beta)$ gives

$$s_1^*(k) \leq \delta_{\infty,6} - 2\eta D^{3+4k}(1-2\beta) + 2D^{4+4k} \leq \delta_{\infty,6},$$

leaving only the periodic elements of $\delta_{k,6}$. ■

5. Proof of Theorem 2 (odd $a \geq 3$)

We first exclude the cases $a = 3, b = 4, 5$ or 6. In all the remaining cases the γ of interest have $t_i \neq a_i$ for almost all i. Now if $t_i = a_i$ infinitely often then from (1.2) we have $M^*(\alpha, \gamma) \leq \eta$. Writing

$$\delta_{\infty,1} = \left(1 - 2\eta + \eta v + \frac{2D}{1-D}\right)\left(1 - \beta + v + \frac{2\beta D}{1-D}\right) = \eta + E_1$$

with

$$E_1 = \eta(a-3)(1-\beta) + \beta D - \frac{2D^2}{1-D} + v\left(1 - \eta + D + \frac{4D^2}{1-D}\right) + \eta v^2 + \frac{4\beta D^2}{(1-D)^2}.$$

When $m \geq 1$ we plainly have $v > 0$ and $E_1 > 0$, when $m = 0$ and $a \geq 5$ we have $E_1 \geq \eta 2(1-\beta) - D > 0$. Hence $\delta_{\infty,1} \geq \eta$ and we can rule out $b_i = a_i - 1$ when $r \geq a+3$ and $b \geq 7$. For $r \leq a+1$ we use that

$$\begin{aligned}\delta_{\infty,3} &= \left(1 - \eta v - \frac{2D}{1-D^2}\right)\left(1 - 3\beta - v - \frac{2\beta D^2}{1-D^2}\right) \\ &> (1 - Dm - 2D)(1 - 3\beta - m\beta + D - m\beta D) \\ &> \eta + E_2 \qquad (5.1)\end{aligned}$$

with

$$E_2 = 1 - \eta - 3\beta - m\beta - D - Dm = \beta\left(m(a-2-\eta) + r\left(1 - \frac{1}{a}\right) - 3 - 2\eta\right).$$

If $m = 0$ and $r = a + 1$ then $E_2 = \beta(a - 1/a - 3 - 2\eta) > 0$ for $a \geq 5$. If $m \geq 1$ and $a \geq 5$ then $E_2 \geq \beta(a - 3 - 2/a - 3\eta) > \beta > 0$. For $a = 3$ ($r = 2$ or 4) we have $E_2 > \beta(\frac{1}{2}m - \frac{8}{3}) > 0$ for $m \geq 6$. One checks numerically that $\delta_{\infty,3} > \eta$ also holds for $b = 3m + 4, m = 1, ..., 5$ and $b = 3m + 2$, $m = 3, 4, 5$, with $\delta_{\infty,7} > \eta$ for $a = 3, b = 8$. Hence we can assume that $t_i \neq a_i$ for almost all i.

We next show that (other than when $r = 2, 4$, $m = 1$) the crucial γ have $t_{2i+1} = \pm 1$. From (1.3) we know that if $|t_{2i+1}| \geq 3$ then $M^*(\alpha, \gamma) < 1 - 3\eta + D$. The rough lower bound

$$\delta_{\infty,1} > 1 - 2\eta - \beta + 3D$$

is enough to rule out this possibility when $r \geq a + 3$. From (5.1) we similarly obtain

$$(1 - 3\eta + D) \leq \delta_{\infty,3} + \beta E_3$$

with

$$E_3 = 3 + 2\eta - m(2 - \eta) - \frac{3r}{a}.$$

Hence if $r = a+1$ we have $E_3 < (2\eta - 3/a) < 0$ and if $m \geq 2$ we have $E_3 < 4\eta - 1 - \frac{3r}{a} < 0$.

Hence we only need consider $t_{2i+1} \neq \pm 1$ when $m = 1$ and $r \leq a - 1$. Now for $m = 1$ if $|t_i| \geq 5$ infinitely often then $M^*(\alpha, \gamma) \leq 1 - 5\beta + D < \delta_{\infty,3} - \beta + 3D < \delta_{\infty,3}$. Hence for $m = 1$, $r \leq a - 1$ we can assume that $t_i = \pm 1, \pm 3$. When $m = 1$, $r \leq a - 1$ we note the rough bound

$$\begin{aligned}\delta_{\infty,5} &> (1 - D + \eta D - D^2)(1 - 4\beta + D - \beta D) \\ &> 1 - 4\beta + 3\beta D.\end{aligned}$$

Now if for example $t_{2i+1} = -3$ and $t_{2i+2} \leq 1$ (likewise if $t_{2i} \leq 1$) then

$$\begin{aligned}s_1^*(2i + 1) &\leq \left(1 - 4\eta + \frac{3D + 3\eta D}{1 - D}\right)\left(1 - \beta + \beta + \frac{3D + 3\beta D}{1 - D}\right) \\ &\leq 1 - 4\eta + \frac{3D + 3\eta D}{1 - D} + (1 - 3\eta)\frac{(3D + 3\beta D)}{1 - D} \\ &< 1 - 4\eta + 6D = 1 - 4\beta + D\left(6 - \frac{4r}{1 + D}\right) < \delta_{\infty,5}.\end{aligned}$$

Hence if $t_{2i+1} = 3\delta$, $\delta = \pm 1$ we can assume that $t_{2i}, t_{2i+2} = -3\delta$. Similarly, if $t_{2i+2} = -3$ and $t_{2i+1} \leq -1$ (likewise for t_{2i+3}) then

$$s_1^*(2i+1) \leq (1-\eta-\eta+3D+3\eta D)(1-\beta-3\beta+3D) < 1-4\beta < \delta_{\infty,5},$$

and we assume that $t_{2i+2} = -3\delta$ implies $t_{2i+1}, t_{2i+3} = \delta$ or 3δ. Now if $t_{2i+1}, t_{2i+2}, t_{2i+3} = -3, 3, -1$ then

$$\begin{aligned} s_2^*(2i+1) &\leq \left(1+\eta-3\eta+\frac{3D-\eta D}{1-D}\right)\left(1+\beta-3\beta+\frac{D(1+\beta)}{1-D}\right) \\ &\leq (1-2\eta+3D)(1-2\beta+D+2\beta D) \\ &< 1-2\beta-2\eta+8D-2\eta D-4\beta D \\ &< \delta_{\infty,5} - 2\beta\frac{(r-4)}{a}. \end{aligned}$$

Hence when $r \geq 4$ we can assume that either $t_{2i+1} = \pm 1$ for almost all i or that $\pm\gamma$ is in S_{-5}. Moreover since

$$\begin{aligned} \delta_{-5} &\leq (1-2\eta+3D)(1-2\beta+3D) \\ &< 1-2\beta-2\eta+10D < \delta_{\infty,5} - 2\beta\left(\frac{r}{a}-5\eta\right), \end{aligned}$$

we do not even need to include δ_{-5} for $r \geq 6$. For $r = 4$ and $5 \leq a \leq 11$ one checks numerically that $\delta_{\infty,5} > \delta_{-5}$. For $r = 4$ and $a \geq 13$ we have

$$\delta_{-5} - \delta_{\infty,5} = \frac{2D}{1-D}\left(1-11\beta-\frac{6D(1-D)}{1+D}+\frac{5D(1-\eta)(1-\beta)}{1-D}\right) > 0,$$

with $\delta_0 > 1-2\eta > \delta_{-5}$.

Suppose now that $b = a+2$. If $a = 5$ or 7 then we can still assume that $t_{2i+1} = \pm 1$ since if $t_{2i+1} = -3$ then

$$s_1^*(2i+1) \leq \left(1-\eta-3\eta+\frac{3D-\eta D}{1-D}\right)\left(1-\beta+\frac{3\beta-D}{1-D}\right) < \delta_{\infty,8}.$$

Hence we assume that $a \geq 9$. We note the lower bound

$$\begin{aligned} \delta_{\infty,9} &\geq (1-2\eta+3D-3\eta D)\,(1-2\beta+D-\beta D) \\ &> 1-2\beta-2\eta+8D-5\eta D-7\beta D+10D^2 > 1-4\beta+4D-12\eta D. \end{aligned}$$

Now if $t_{2i+2} = -3\delta$, $\delta = \pm 1$, then we can assume that at least one of t_{2i+1}, $t_{2i+3} = 3\delta$, since if $t_{2i+2} = -3$, $t_{2i+1} = t_{2i+3} = 1$ then

$$\begin{aligned} s_1^*(2i+1) &\le \left(1 - \eta + \frac{\eta + D}{1-D}\right)\left(1 - \beta - 3\beta + \frac{D + \beta D}{1-D}\right) \\ &< 1 - 4\beta + \frac{D+\beta D}{1-D} + \frac{D+\eta D}{1-D}(1-3\beta) \\ &< 1 - 4\beta + 2D < \delta_{\infty,9}. \end{aligned}$$

Similarly if $t_{2i+2} = -3\delta$ and $t_{2i+1} = \delta$ then $t_{2i} = \delta$, since if $t_{2i+2}, t_{2i+1} = -3, 1$, $t_{2i} \le -1$ then

$$\begin{aligned} s_1^*(2i+1) &\le \left(1 - \eta + \eta - D + \frac{\eta D + D^2}{1-D}\right)\left(1 - \beta - 3\beta + \frac{3D - 3\beta D}{1-D}\right) \\ &< (1 - D + 2\eta D)(1 - 4\beta + 3D - 2\beta D) \\ &< 1 - 4\beta + 2D + 2\beta D + 2\eta D < \delta_{\infty,9}. \end{aligned}$$

(Likewise $t_{2i+4} = \delta$ if $t_{2i+2} = -3\delta$ and $t_{2i+3} = \delta$). Also if $t_{2i+1} = \delta$ and $t_{2i+1+\delta'} = -\delta$ with δ and $\delta' = \pm 1$ then $t_{2i+1-\delta'} = -\delta$ or -3δ, since if $t_{2i+1}, t_{2i+2} = -1, 1$, $t_{2i} \le -1$ then

$$\begin{aligned} s_1^*(2i) &\le \left(1 - \beta - \beta + \frac{D + \beta D}{1-D}\right)\left(1 - \eta - \eta + \frac{D + \eta D}{1-D}\right) \\ &< (1-2\beta)(1-2\eta) + \frac{D}{1-D}((1+\beta)(1-\eta) + (1+\eta)(1-2\beta)) \\ &< 1 - 2\beta - 2\eta + 6D < \delta_{\infty,9}. \end{aligned}$$

So if $t_{2i+1} = -1$, t_{2i+2} (or t_{2i}) $= -1$ infinitely often then $s_1^*(2i+1)$ is at most

$$\left(1 - \eta - \eta + 3D - 3\eta D + 3D^2 - \eta D^2 + \frac{-D^3 + \eta D^3}{1-D}\right)\left(1 - \beta - \beta + \frac{D - D\beta}{1-D}\right).$$

Thus if $\pm\gamma$ has $t_{2i} = -3$ infinitely often and $\pm\gamma$ is not in S_{-5} we must have infinitely many blocks t_{2i+1}, t_{2i+2} or t_{2i-1}, t_{2i-2} of the form $1, 1$ and hence $M^*(\alpha, \gamma) \le \delta_{\infty,9}$. This only leaves the γ with $t_i = \pm 1$, for almost all i with $t_{2i+1}, t_{2i+1\pm 1}$ of the form $-1, -1$ or $1, 1$ at most finitely often; which leaves just the $\pm\gamma$ in S_0. For $a \ge 11$ we have

$$\delta_{-5} > \delta_{\infty,9} - 2\eta D^2 + 2D(1-\beta)(1-2\eta) > \delta_{\infty,9}.$$

One checks numerically that this is also true for $a = 9$.

Thus in the remaining cases we can assume that $t_{2i+1} = \pm 1$ for all i. Suppose that γ and $1 - \alpha - \gamma$ satisfy

$$t_{2i+1} = -1 \Rightarrow (m - S) \le t_{2i}, t_{2i+2} \le (m + 2 + R) \tag{5.2}$$
$$t_{2i+1} = 1 \Rightarrow -(m + 2 + R) \le t_{2i}, t_{2i+2} \le -(m - S)$$

for almost all i, with R and S minimal.

Suppose first that $R \ge S$ and that γ (or $1-\alpha-\gamma$) has $t_{2i+1} = 1$, $t_{2i+2} = -(m+2+R)$ infinitely often. Hence

$$d^-_{2i+1} \le \eta + \frac{(-m + S)D + \eta D}{1 - D}$$
$$d^+_{2i+1} \le -(m + 2 + R)\beta + \frac{D}{1 - D} + \frac{(-m + S)\beta D}{1 - D},$$

and

$$\begin{aligned} s^*_1(2i + 1) &\le \left(1 - \eta v + \frac{RD}{1 - D}\right)\left(1 - 3\beta - v - \frac{\beta R(1 - 2D)}{1 - D}\right) \\ &= \delta_{\infty,5} - \frac{R\beta}{1 - D}\left(1 - \eta + D + 2\eta D v + \frac{RD(1 - 2D)}{1 - D}\right) \\ &< \delta_{\infty,5} - \frac{2\beta}{1 - D}(1 - \eta + D), \end{aligned}$$

for $R \ge 2$. Since

$$\begin{aligned} \delta_{\infty,5} &= \delta_{\infty,3} + \frac{2\beta/a}{1 - D}\left(1 - 3\beta + D + \beta D - \frac{m\beta(1 + D^2)}{1 - D} + \frac{D^2(1 - 2/b + D)}{1 - D}\right) \\ &< \delta_{\infty,3} + \frac{2\beta/a}{1 - D}, \end{aligned}$$

and

$$\begin{aligned} \delta_{\infty,5} &= \delta_{\infty,1} - \frac{2\beta}{1 - D}\left(1 - \frac{r}{a}(1 - D) + \eta - D - \frac{2\eta D}{1 - D} + \frac{2(m + 1)D^2}{1 - D}\right) \\ &< \delta_{\infty,1} - \frac{2\beta}{1 - D}(-1 + \eta + D - 2\eta D), \end{aligned}$$

we immediately obtain $M^*(\alpha, \gamma) < \delta_{\infty,3}$ and $\delta_{\infty,1}$ for $R \ge 2$. Hence we can assume that $S \le R \le 0$ and (since plainly $R + S \ge -2$) the only possibilities are $R = S = 0$ or $R = 0, S = -2$.

Similarly suppose that $R < S$ and $t_{2i+1} = -1$, $t_{2i+2} = (m - S)$. Then

$$d^-_{2i+1} \le -\eta + (m + 2 + R)D + \frac{\eta D}{1-D} + \frac{(-m+S)D^2}{1-D},$$

$$d^+_{2i+1} \le (m-S)\beta + \frac{D}{1-D} + \frac{(-m+S)\beta D}{1-D}, \tag{5.3}$$

bounding $s^*_1(2i+1)$ by

$$\begin{aligned}&\left(1 - 2\eta + \eta v + \frac{2D}{1-D} + \frac{D}{1-D}(S - 2 + 2\eta - 2mD)\right)\\ &\qquad\left(1 - \beta + v + \frac{2\beta D}{1-D} - \frac{\beta}{1-D}(S(1-2D) + 2D + 2mD - 2\eta)\right)\\ &= \delta_{\infty,1} - \frac{2D}{1-D}\left(\frac{1}{2}S(a-3) + \frac{1}{2}(S-2)U_1 + U_2\right), \end{aligned} \tag{5.4}$$

where (since $m\beta < \eta$)

$$\begin{aligned}U_1 &= S\frac{\beta(1-2D)}{1-D} + \frac{2D}{1-D}\left(2 - \beta - 3D - m\beta(1-2D)\right) > 0,\\ U_2 &= \frac{2D(1-3D)}{1-D} + \eta + \frac{4D^2(m\beta)}{1-D} + (m\beta)(2 - 3\eta - D) + 2(m\beta)^2\eta > 0.\end{aligned}$$

Hence $M^*(\alpha,\gamma) < \delta_{\infty,1}$ for $S \ge 2$. For $r = a + 1$ we have

$$\delta_{\infty,1} - \delta_{\infty,3} = \frac{2D}{1-D}W,$$

with

$$\begin{aligned}W &= 1 - \eta - 2\beta + D + \frac{2\eta D}{1-D^2} + \frac{4\beta D}{1+D} - \frac{2D^2(3 + D - \beta D - D^2)}{(1+D)(1-D^2)}\\ &< 1 - \eta + 2\eta D - \frac{4}{3}\beta.\end{aligned}$$

When $a \ge 5$ the bound $\frac{1}{2}S(a-3) \ge 2$ gives $M^*(\alpha,\gamma) < \delta_{\infty,3}$ for $S \ge 2$. When $a = 3$ we have

$$\begin{aligned}U_2 &= 1 - \eta + D - 2\eta D + \frac{2D(1-3D)}{1-D} - \frac{r}{3}\beta(1-2D) + 2\eta(m\beta)^2 + \frac{4D^2(m\beta)}{1-D}\\ &> 1 - \eta + 3D - 2\eta D - \frac{4}{3}\beta > W\end{aligned}$$

and still obtain $M^*(\alpha,\gamma) < \delta_{\infty,3}$. Writing

$$\delta_{\infty,1} = \delta_{\infty,5} + \frac{2D}{1-D}W_1,$$

where

$$W_1 = (a-r+1) - 2\beta\left(1-\frac{r}{a}\right) - \frac{2D}{1-D}\left(1-\eta-\beta\left(1-\frac{r}{a}\right)\right),$$

it is clear that for $4 \le r \le a-1$ the rough bounds $W_1 < (a-3)$ and $\frac{1}{2}S(a-3) \ge (a-3)$ are enough to give $M^*(\alpha,\gamma) < \delta_{\infty,5}$ when $S \ge 2$. Hence when $r \ge 4$ we must have $S \le 0$, so we can assume that

$$t_{2i+1} = \delta \Rightarrow t_{2i+1\pm1} = -m\delta, \text{ or} - (m+2)\delta,$$

for $\delta = 1$ or -1, so that the coefficients in the expansion of γ consists of blocks B_m, B_n, B'_m or B'_n.

When $r = 2$ and $a \ge 5$ the bounds $W_1 < (a-1)$ and $\frac{1}{2}S(a-3) \ge 2a-6$ give $M^*(\alpha,\gamma) < \delta_{\infty,5}$ for $S \ge 4$. Thus we can assume that $S \le 2$ and hence that

$$t_{2i+1} = \delta \Rightarrow t_{2i+1\pm1} = -(m-2)\delta, -m\delta, \text{ or} - (m+2)\delta, \tag{5.5}$$

With a bit more work we show that this is also true for $a = 3, b = 3m+2, m \ge 2$: For $a = 3$, $r = 2$ observe that $(m\beta) = \eta - \frac{2}{3}\beta$ and for $S \ge m$

$$U_1 \ge \eta + \frac{D}{1-D}\left(1 - \frac{5}{9}\beta + \frac{4}{3}D^2 - \frac{41}{9}\beta D\right) > \eta$$

and

$$\begin{aligned} U_2 &= 1-\eta+D+\frac{2D}{1-D}((1-D)^2-\eta-\frac{4}{3}\beta D) - \frac{2}{3}\beta(1-2D) + 2\eta(m\beta)^2 \\ &> 1-\eta+D-\frac{2}{3}\beta > -4\eta+2-\frac{2}{3}\beta \end{aligned}$$

while

$$W_1 = 2 - \frac{2}{3}\beta - \frac{2D}{1-D}(1-\eta-\frac{1}{3}\beta) < 2 - \frac{2}{3}\beta.$$

Hence $M^*(\alpha,\gamma) < \delta_{\infty,5}$ for $\frac{1}{2}(S-2) \ge 4$. Thus it remains to consider $S < m$ or the odd cases $S = 8, m \le 8$, $S = 6, m \le 6$ or $S = 4, m \le 4$. Notice that $t_{2i} = (m+2+R)$

infinitely often implies that $M^*(\alpha,\gamma) \le 1-(m+2+R)\beta+D$, so that we can assume that $(m+2+R) < (1+D-\delta_{\infty,5})/\beta$ is at most $12,11,10,7,6,5,4$ as $m = 8,7,6,5,4,3,2$. Using this bound for $(m+2+R)$ in (5.3) gives numerically $M^*(\alpha,\gamma) < \delta_{\infty,5}$ for the oddments. Thus it remains only to deal with $4 \le S < m$, $R < S$. In particular (5.2) implies that the t_i are alternately greater or less than 0, so that the t_{2i+1} are constant, improving (5.4) to

$$\begin{aligned} M^*(\alpha,\gamma) &\le \left(1-2\eta+\eta v+\frac{(R+2)D}{1-D}\right)\left(1-\beta+v-S\beta+\frac{(R+2)\beta D}{1-D}\right) \\ &= \delta_{\infty,5}+DW_2+D(R+2-S)v, \end{aligned}$$

with

$$\begin{aligned} W_2 &= 2-\frac{2}{3}\beta-2v-S(1-\beta)+\frac{(R+2)}{1-D}\left(1-2D+2Dv-S\beta+\frac{D\beta(R+2)}{1-D}\right) \\ &< (4+R-S)(1-\beta). \end{aligned}$$

Hence $M^*(\alpha,\gamma) < \delta_{\infty,5}$ for $R \le S-4$, so we can assume that $S \ge 4$ and $R = (S-2) \ge \frac{1}{2}S$. But $t_{2i+1} = 1$, $t_{2i+2} = -(m+2+R)$ then gives

$$\begin{aligned} s_1^*(2i+1) &\le \left(1-\eta v+\frac{SD}{1-D}\right)\left(1-3\beta-v-\frac{1}{2}S\beta+\frac{S\beta D}{1-D}\right) \\ &= \delta_{\infty,5}-\frac{1}{2}\frac{SD}{1-D}\left((3-4\beta)\left(1-\eta v+\frac{SD}{1-D}\right)-2(1-3\beta-v)\right) \\ &< \delta_{\infty,5}. \end{aligned}$$

So we can assume that $S \le 2$ and (5.5) holds.

Suppose now that $r \ge a+3$ with $m \ge 1$. From the previous discussion we can assume that $t_{2i+1} = -1$ and $t_{2i+2} = m$ or $(m+2)$ for almost all i.

Now if $t_{2i+2} = m$ then

$$\begin{aligned} s_1^*(2i+1) &\le \left(1-\eta-\frac{\eta}{1-D}+\frac{(m+2)D}{1-D}\right)\left(1-\beta+m\beta-\frac{D}{1-D}+\frac{(m+2)\beta D}{1-D}\right) \\ &= \delta_{\infty,1}. \end{aligned}$$

Hence either $\pm\gamma$ is in S_{-2} or $M^*(\alpha,\gamma) \le \delta_{\infty,1}$. For $r \ge a+3$,

$$\delta_{-2} = \delta_{\infty,1}+2\beta\left(\frac{r-(a+2)}{a}+\eta v+\frac{2D}{1-D}-\frac{2D}{a}\right) > \delta_{\infty,1}.$$

Similarly suppose that $4 \le r \le a-1$ and that γ is not in S_{-5} when $r = 4$ and $a \ge 13$. Then from the above we can assume that γ has $t_{2i+1} = 1$, $t_{2i+2} = -m$ or $-(m+2)$ for almost all i. If $t_{2i+1} = -(m+2)$ then

$$s_1^*(2i+1) \le \left(1 - \beta - (m+2)\beta + \frac{D - m\beta D}{1-D}\right)\left(1 - \eta + \frac{\eta - mD}{1-D}\right) = \delta_{\infty,5}.$$

Hence either $\pm\gamma$ is in S_0 or $M^*(\alpha,\gamma) \le \delta_{\infty,5}$. Since $v \le \eta$ and $r \le a-1$,

$$\delta_0 = \delta_{\infty,5} + 2\beta\left(\frac{a-r}{a} - \eta v\right) > \delta_{\infty,5}.$$

Suppose that $r = a+1$ with $m \ge 1$ then again we can assume that $t_{2i+1} = 1$, $t_{2i+2} = -m$ or $-(m+2)$ for almost all i. If $t_{2i+2} = t_{2i} = -m$ then $s_3^*(2i+1)$ is at most

$$\left(1 - \eta - \frac{\eta}{1-D} + mD + \frac{(m+2)D^2}{1-D}\right)\left(1 - \beta + m\beta - \frac{D}{1-D} + \frac{(m+2)\beta D}{1-D}\right)$$
$$= \delta_{-2} - \frac{2\beta D(1+3D)}{1-D} + \frac{2\beta D}{a} - 4Dv < \delta_{-2} < \delta_{\infty,3}.$$

Hence we can assume that the expansion of γ eventually consists of B_ms and B_ns with no consecutive B_ms. Thus either the B_ms and B_ns eventually alternate and γ is in S_{-1} or γ contains infinitely many consecutive B_ns, $t_{2i+1\pm1} = -(m+2)$ and

$$d^+_{2i+1} \le -(m+2)\beta + \frac{D}{1-D} - \frac{m\beta D}{1-D^2} - \frac{(m+2)\beta D^2}{1-D^2}$$
$$d^-_{2i+1} \le \frac{\eta}{1-D} - \frac{(m+2)D}{1-D^2} - \frac{mD^2}{1-D^2}$$

giving $s_1^*(2i+1) \le \delta_{\infty,3}$. Clearly $\delta_{-1} > \delta_{\infty,3}$ for $r = a+1$.

It remains to consider $m = 0$ or $r = 2$, $m \ge 2$.

Suppose first that $m = 0$ and $r \ge a+3$. By the previous analysis we can assume that $t_{2i+1} = \delta_i$ and $t_{2i+1\pm1} = 0$ or $-2\delta_i$ with $\delta_i = \pm 1$. Hence if $\pm\gamma$ is not in S_{-2} we must have $t_{2i+2} = 0$ infinitely often. Now if $t_{2i+2} = 0$ we can assume that $t_{2i+1} \ne t_{2i+3}$, since $t_{2i+1} = t_{2i+3} = -1$ would give

$$s_1^*(2i+2) \le \left(1 - \eta - \eta + \frac{2D - \eta D}{1-D}\right)\left(1 - \beta - D + \frac{2\beta D - D^2}{1-D}\right) < \delta_{\infty,2}.$$

We assume that $t_{2i+1}, t_{2i+2}, t_{2i+3} = 1, 0, -1$ giving

$$s_1^*(2i+2) = (1-\eta-\eta+\eta d_{2i+3}^+)(1-\beta+D+Dd_{2i}^-)$$
$$s_3^*(2i+1) = (1-\eta-\eta-\eta d_{2i}^-)(1-\beta+D-Dd_{2i+3}^+).$$

Hence, writing $d^* = \max\{d_{2i+3}^+, -d_{2i}^-\}$ we plainly have

$$\min\{s_1^*(2i+2), s_3^*(2i+1)\} < (1-2\eta+\eta d^*)(1-\beta+D-Dd^*) \le d_{\infty,2}^*,$$

since $d^* \le (2\beta - D)/(1-D)$. The claim follows on checking that

$$\delta_{-2} = \delta_{\infty,2} + \frac{2\beta}{(1-D)^2}\left(\frac{r-(a+3)}{a} + D\left(6 - \frac{3}{a} - 2\eta + \eta D - D\right)\right) > \delta_{\infty,2}.$$

Suppose that $b = a+1$, $a \ge 5$. We note the rough bound

$$\delta_{\infty,4} > (1-\eta D)(1-3\beta+D) > 1-3\beta+D-\eta D.$$

From above we have $t_{2i+1} = \delta_i$, $t_{2i+1\pm1} = 0$ or $-2\delta_i$, $\delta_i = \pm 1$ for almost all i. If $t_{2i+1\pm1} = -2$ then

$$\begin{aligned} s_1^*(2i+1) &\le \left(1-\eta+\eta-2D+\frac{\eta D}{1-D}\right)\left(1-\beta-2\beta+\frac{D}{1-D}\right) \\ &\le 1-3\beta+D-\eta D-2D(1-\eta-3\beta) < \delta_{\infty,4}. \end{aligned}$$

So we can assume that γ does contain consecutive $t_{2i} = t_{2i+2} = -2\delta_i$. Similarly we rule out the possibility of $t_{2i} = t_{2i+2} = 0$ infinitely often when γ is not in S_{-4}, since otherwise we can assume that $\pm\gamma$ has blocks $t_{2i+1} = -1$, $t_{2i} = t_{2i+2} = 0$ and either $t_{2i+3} = -1$ and

$$\begin{aligned} s_1^*(2i+1) &\le \left(1-\eta-\eta+\frac{\eta D}{1-D}\right)\left(1-\beta-D+\frac{2\beta D - D^2}{1-D}\right) \\ &< 1-2\eta-\beta+D+3\eta D+2\beta D = 1-3\beta-D+5\eta D < \delta_{\infty,4}, \end{aligned}$$

or $t_{2i+3} = 1$ and $t_{2i+4} = -2$ and

$$s_1^*(2i+3) \le \left(1-\eta+\eta-\eta D+\frac{\eta D^2}{1-D}\right)\left(1-\beta-\frac{2\beta}{1-D^2}+\frac{D}{1-D}\right) < \delta_{\infty,4}.$$

Hence apart from S_{-4} it is enough to consider γ such that the t_{2i} alternate between 0 and $-2\delta_i$. Thus if $\pm\gamma$ is not in S_{-1} we can assume that γ contains infinitely many blocks $t_{2i-1}, ..., t_{2i+5} = 1, -2, 1, 0, -1, 2, -1$. Hence we have

$$s_1^*(2i) = (1 - \beta - 2\beta + D + Dd_{2i-2}^-)(1 - \eta + \eta - \eta D + 2D^2 - \eta D^2 + \eta D^2 d_{2i+5}^+)$$
$$s_3^*(2i+3) = (1 - \beta - 2\beta + D - Dd_{2i+5}^+)(1 - \eta + \eta - \eta D + 2D^2 - \eta D^2 - \eta D^2 d_{2i-2}^-)$$

and, setting $d^* = \max\{d_{2i-2}^-, -d_{2i+5}^+\}$,

$$\min\{s_1^*(2i), s_3^*(2i+3)\} \le (1 - 3\beta + D + Dd^*)(1 - \eta D + 2D^2 - \eta D^2 - \eta D^2 d^*) \le \delta_{\infty,4},$$

since $d^* \le D/(1-D) - 2\beta D/(1-D^2)$. Hence $M^*(\alpha, \gamma) \le \delta_{\infty,4}$ for $\pm\gamma$ not in S_{-1} or S_{-4} where

$$\delta_{-4} = \delta_{\infty,4} - \frac{2D^2}{(1-D^2)^2}\left(3 - 3\beta - 3\eta + 4D - 2D^2 + \beta D^2 + \eta D^2 - D^3\right) < \delta_{\infty,4}$$

and plainly $\delta_{-1} > \delta_{\infty,4}$.

We are left to deal with $r = 2$, $m \ge 2$.

Suppose first that $b = ma + 2$ with $m \ge 3$. Then from above we can assume that the expansion of γ has $t_{2i+1} = 1$, $t_{2i+2} = -(m-2), -m$ or $-(m+2)$. Now if $t_{2i+2} = -(m-2)$ and t_{2i} (or t_{2i+4}) $\ge -m$ then $s_3^*(2i+1)$ is bounded by

$$\left(1 - 2\eta + mD - \frac{\eta D}{1-D} + \frac{(m+2)D^2}{1-D}\right)\left(1 - \beta + (m-2)\beta - \frac{D}{1-D} + \frac{(m+2)\beta D}{1-D}\right)$$
$$= \delta_{\infty,6} - \frac{2Dv}{(1-D^2)}(2 - 3D - 3D^2) + \frac{2D^2(1-3D)}{1-D^2} - \frac{2\beta D^2(1-5D^2)}{(1-D)(1-D^2)} < \delta_{\infty,6}.$$

Hence we assume that $t_{2i+2} = -(m-2)$ implies that $t_{2i+4}, t_{2i} = -(m+2)$. So if $\pm\gamma$ is not in S_0 or S_{-3} we can assume that $t_{2i} = -(m+2)$, $t_{2i+2} \le -m$ infinitely often and

$$s_1^*(2i) \le \left(1 - \beta - \frac{(m+2)\beta}{1-D^2} + \frac{D}{1-D} - \frac{(m-2)\beta D}{1-D^2}\right)\left(1 + \frac{\eta D}{1-D} - \frac{mD}{1-D}\right)$$
$$= \delta_{\infty,6}.$$

Checking

$$\delta_0 = \delta_{-3} + \frac{2}{b}\left(1 - 3\eta + D + \frac{2\eta}{b}\right) > \delta_{-3},$$

and

$$\delta_{-3} = \delta_{\infty,6} + \frac{2D}{1+D}\left(1 - 3\beta - v + \frac{2\beta D}{1+D}\right) > \delta_{\infty,6}$$

gives the claimed result.

Suppose that $b = 2a+2$. We can assume here that $t_{2i+1} = \delta_i$, $t_{2i+1\pm 1} = 0, -2\delta_i$ or $-4\delta_i$ with $\delta_i = \pm 1$. We note the lower bound

$$\delta_{\infty,7} \geq 1 - 5\beta + D - \eta D.$$

If the expansion contains blocks of the form $t_{2i} = 0$, $t_{2i+1} = -1$, $t_{2i+2} \neq 4$ then

$$\begin{aligned} s_1^*(2i) &\leq \left(1 - \beta + \frac{D}{1-D}\right)\left(1 - \eta - \eta + 2D - \frac{\eta D}{1-D} + \frac{4D^2}{1-D}\right) \\ &< 1 - 2\eta - \beta + 5D - 2\beta D - 3\eta D + 8D^2 \\ &= 1 - 5\beta + D - \eta D - 6\beta D + 8D^2 < \delta_{\infty,7}. \end{aligned}$$

Hence we assume $t_{2i} = 0$ implies that $t_{2i-2}, t_{2i+2} = \pm 4$. Similarly if $t_{2i+2} = -4$, $t_{2i+1} = 1$, and $t_{2i} \leq -2$ then

$$\begin{aligned} s_1^*(2i+1) &\leq \left(1 - \eta + \eta - 2D + \frac{\eta D}{1-D}\right)\left(1 - \beta - 4\beta + \frac{D}{1-D}\right) \\ &< 1 - 5\beta - D + \eta D + 10\beta D - 4D^2 \\ &= 1 - 5\beta + D - \eta D - 2\beta D\left(2a - \frac{2}{a} + \eta - 5\right) < \delta_{\infty,7}. \end{aligned}$$

So we may assume that $t_{2i} = 4\delta_i$ implies that $t_{2i-2}, t_{2i+2} = 0$. Hence we need only consider $\pm\gamma$ in S_0 or γ where the t_{2i} are alternately 0 and $-4\delta_i$. Of these γ either $\pm\gamma$ is in S_{-3} or $\pm\gamma$ has infinitely many blocks $t_{2i}, ..., t_{2i+4} = -4, 1, 0, -1, 4$, giving

$$\begin{aligned} s_1^*(2i) &= \left(1 - \beta + d_{2i}^-\right)\left(1 - \eta + \eta - \eta D + \eta D d_{2i+3}^+\right) \\ s_3^*(2i+3) &= \left(1 - \beta - d_{2i+3}^+\right)\left(1 - \eta + \eta - \eta D - \eta D d_{2i}^-\right) \end{aligned}$$

and, setting $d^* = \max\{d_{2i}^-, -d_{2i+3}^+\} \leq -\frac{4\beta}{1-D^2} + \frac{D}{1-D}$ we obtain

$$\min\{s_1^*(2i), s_3^*(2i+3)\} \leq (1 - \beta + d^*)(1 - \eta D - \eta D d^*) \leq \delta_{\infty,7}.$$

Finally, $\delta_0 > \delta_{-3}$ just as for $m \geq 2$ and $\delta_{-3} > \delta_{\infty,7}$ since

$$\delta_{-3} - \delta_{\infty,7} = \frac{2\beta D}{(1-D)}\left(2 - 7\eta + \frac{2D}{(1+D)(1-D^2)}(11 + 2D - D^2 - 3\eta D(1+D))\right).$$

It remains only to deal with $(a,b) = (3,4), (3,5), (3,6), (5,7), (7,9)$.

For $(5,7)$ and $(7,9)$ we know from above that $t_{2i+1} = \delta_i$, $\delta_i = \pm 1$, $t_{2i+1\pm 1} = \pm\delta_i$ or $-3\delta_i$. Now if $t_{2i} = -3\delta_i$ then we can assume that $t_{2i+2} = t_{2i-2} = \delta_i$, since if for example $t_{2i} = -3$, $t_{2i+1} = t_{2i-1} = 1$, $t_{2i+2} \leq -1$ then

$$s_1^*(2i) \leq \left(1 - \beta - 3\beta + \frac{D + \beta D}{1-D}\right)\left(1 - \eta + \eta - D + \frac{\eta D + D^2}{1-D}\right) < \delta_{\infty,8}.$$

Also, if $t_{2i-1} = t_{2i} = t_{2i+1} = -\delta$, $\delta = \pm 1$, then $t_{2i-2} = t_{2i+2} = 3\delta$, since if $t_{2i+1} = t_{2i} = t_{2i-1} = -1$ and $t_{2i-2} \leq 1$ then $s_1^*(2i-1)$ is at most

$$\left(1 - 2\eta + \frac{D + \eta D}{1-D}\right)\left(1 - 2\beta - D + 3\beta D - D^2 - \beta D^2 + \frac{D^3 + \beta D^3}{1-D}\right) < \delta_{\infty,8}.$$

Now if $t_{2i+1} = -\delta$ we can assume that we do not have $t_{2i} = t_{2i+2} = -\delta$ since $t_{2i} = t_{2i+1} = t_{2i+2} = -1$ would imply

$$s_1^*(2i) \leq \left(1 - \beta - \beta + \frac{D + \beta D}{1-D}\right)\left(1 - \eta - \eta - D + \frac{\eta D + D^2}{1-D}\right) < \delta_{\infty,8}.$$

Similarly if $t_{2i+1} = \delta$, $t_{2i+1\pm 1} = -\delta$ and t_{2i-1} (or t_{2i+3}) $= -\delta$ then t_{2i-2} (or t_{2i+4}) $= 3\delta$, since for example $t_{2i-1}, t_{2i}, t_{2i+1}, t_{2i+2} = -1, -1, 1, -1$ and $t_{2i-2} \leq 1$ then $s_1^*(2i-1)$ is at most

$$\left(1 - 2\eta + D + \eta D - D^2 + \frac{\eta D^2 + D^3}{1-D}\right)\left(1 - 2\beta + D - \beta D + \frac{D^2 + \beta D^2}{1-D}\right) < \delta_{\infty,8}.$$

With these exclusions we readily find that $M^*(\alpha,\gamma) < \delta_{\infty,8}$ if the t_{2i+1} are eventually constant and $\pm\gamma$ is not in S_0 or S_{-3} and if the t_{2i+1} eventually alternate and $\pm\gamma$ is not in S_{-4}. Hence we may assume that $\pm\gamma$ has infinitely many blocks of the form $t_{2i-1} = t_{2i+1} = 1$ and $t_{2i+3} = -1$. Now $t_{2i} = -1$ or -3. If $t_{2i} = -3$ then $s_1^*(2i)$ is at most

$$\left(1 - 4\beta + \frac{D + \beta D + D^2 - 3\beta D^2}{1-D^2}\right)\left(1 + D - \eta D + \frac{D^2 + \eta D^2 - 3D^3 + \eta D^3}{1-D^2}\right)$$

which equals $\delta_{\infty,8}$. So we assume that $t_{2i} = -1$. If $t_{2i+2} = -1$ then $t_{2i+4} = 3$ and

$$s_3^*(2i+3) \le \left(1 + D - \eta D + D^2\right)\left(1 - 4\beta + \frac{D + \beta D + D^2 - 3\beta D^2}{1 - D^2}\right) < \delta_{\infty,8}.$$

If $t_{2i+2} = 1$ then we can assume that $t_{2i+4} = -1$. But if $M^*(\alpha,\gamma) > \delta_{\infty,8}$ then the previous exclusions force the subsequent coefficients $t_{2i+1+2j}$ to alternate between ± 1 and the $t_{2i+2j+2}$ between ± 1 so that γ is in S_{-4}. One checks numerically that $\delta_0 > \delta_{-3} > \delta_{-4} > \delta_{\infty,8}$.

Finally we deal with $a = 3$, $b = 4, 5, 6$.

In all cases we can rule out $t_{2i} = b$ since $\beta < \delta_{\infty,i}$, $i = 10, 11$ or 12.

We begin with $(a, b) = (3, 4)$. Suppose that $t_{2i+1} = a$ then

$$s_2^*(2i) = \eta(1 + \beta + d_{2i}^-)(1 - \beta - d_{2i+1}^+),$$
$$s_4^*(2i+1) = \eta(1 + \beta + d_{2i+1}^+)(1 - \beta - d_{2i}^-),$$

and if $d_{2i+1}^+ \ge aD - \beta D(2+\eta)/(1-D)$ the minimum of these is certainly at most $\eta(1 - (\beta + d_{2i+1}^+)^2) < \delta_{\infty,10}$. Hence we can assume that $t_{2i+2}, t_{2i+3}, \ldots$ (and likewise $t_{2i}, t_{2i-1}, \ldots$) must take the form $-2, \ldots$ or $0, \pm 1, \ldots$. Also we can successively rule out blocks t_{2i+1}, t_{2i+2}, of the form $-1, -2$ (and hence also blocks $a, 0, 1$, since its negative is $a, -2, -1$) or $-1, 0, -1, 0$ or $1, -2, 1$ (or their 'negatives') else

$$s_1^*(2i+1) \le \left(1 - 2\eta + \frac{2D + \eta D}{1 - D}\right)\left(1 - 3\beta + \frac{aD}{1-D}\right) < \delta_{\infty,10},$$

or

$$s_1^*(2i+2) \le (1 - 2\eta + aD\eta)\left(1 - \beta - D + \frac{2\beta D - D^2}{1 - D}\right) < \delta_{\infty,10},$$

or

$$s_1^*(2i+2) \le \left(1 - 3\beta + D + D^2\right)(1 + \eta D) < \delta_{\infty,10}.$$

In particular $t_{2i+1} = -1$ forces $t_{2i} = t_{2i+2} = 0$. These also rule out blocks $-1, 0, -1$ and we can further dismiss $t_{2i-1}, t_{2i}, t_{2i+1} = 1, 0, -1$ since

$$s_1^*(2i+1) = \eta(1 - \beta + D - D(-d_{2i-2}^-))(1 - \beta + d_{2i+1}^+)$$
$$s_3^*(2i-2) = \eta(1 - \beta + D - Dd_{2i+1}^+)(1 - \beta + (-d_{2i-2}^-))$$

and since $d^+_{2i+2} \leq (aD - D^2)/(1 - D^2)$ the minimum of these is certainly at most $\eta(1 - \beta + d^+_{2k+1})(1 - \beta + D - Dd^+_{2k+1}) \leq \delta_{\infty,10}$. In particular this rules out blocks $1, 0$. Hence we can assume that the sequence for γ consists entirely of blocks $F_2 = (a, -2)$, $M = (a, -2, 1, -2)$ or $M' = (a, 0, -1, 0)$. We suppose that γ does not have period F_2 or M or M'. If a block $t_{2i+1}, t_{2i+2}, ... = M, ...$ is preceeded by blocks M' or $M'F_2$ or F_2F_2 then

$$d^+_{2i+1} \leq -2\beta + D - 2\beta D + \frac{aD^2 - D^3}{1 - D^2},$$
$$d^-_{2i} \geq -2\beta + aD - 2\beta D + \frac{aD^2 - 2\beta D^2 + D^3 - 2\beta D^3}{1 - D^2}$$

and $s^*_4(2i + 1) = \eta(1 + \beta + d^+_{2i+1})(1 - \beta - d^-_{2i}) < \delta_{\infty,10}$. Hence we can assume that the sequence consists of blocks of M's separated by single F_2's (or its negative with M''s replacing the M's). Suppose that we have a block $...|t_{2i+1}, t_{2i+2}, ...$ of the form $...F_2M^sF_2|M^tF_2, ...$ with $t \geq s + 1$ then

$$d^+_{2i+1} \leq \frac{-2\beta + D - 2\beta D + aD^2}{1 - D^2} + (a - 1)D^{1+2t},$$

$$d^-_{2i} \geq -2\beta + \frac{aD - 2\beta D + D^2 - 2\beta D^2}{1 - D^2} + (a - 1)D^{2+2s}(1 - D)$$

and $s^*_4(2i + 1) \leq \delta_{\infty,10} + \eta(a - 1)D^{2s+2}(D(1 + \beta D) - (1 - \beta)(1 - D)) < \delta_{\infty,10}$. This leaves only the periodic elements of $S_{k,10}$.

Suppose next that $(a, b) = (3, 5)$. If $t_{2i+1} = a$ then in the same way we can rule out $d^+_{2i+1} \geq (\beta - D)/(1 - D)$ or $d^+_{2i+1} \leq -3\beta + (D - \beta D)$ and deduce that $t_{2i+2}, ...$ (and $t_{2i}, ...$) take the form $1, -1, ...$ or $-1, ...$ or $-3, a, ...$ or $-3, 1, ...$ We can also successively rule out blocks t_{2i}, t_{2i+1}, of the form $-3, -1$ and $t_{2i-1}, t_{2i}, ... = 1, -3, 1, ...$ else

$$s^*_1(2i) \leq \left(1 - 4\beta + \frac{aD + \beta D}{1 - D}\right)\left(1 - 2\eta + \frac{3D + \eta D}{1 - D}\right) < \delta_{\infty,11}$$

or

$$s^*_1(2i) \leq \left(1 - 4\beta + \frac{D + \beta D}{1 - D}\right)\left(1 + \frac{D + \eta D}{1 - D}\right) < \delta_{\infty,11}.$$

In particular there are no $t_{2i} = 3$ and the $t_{2i} = -3$ are confined to blocks $a, -3, a$ or $a, -3, 1$ or $1, -3, a$. Likewise we can rule out blocks $t_{2i}, t_{2i+1} = -1, -1$ with $t_{2i+2} = -1$ or with $t_{2i-1} \leq 1$ else

$$s_1^*(2i) \leq (1 - 2\beta + aD + \beta D)(1 - 2\eta - D + a\eta D + D^2) < \delta_{\infty,11}$$

or

$$s_1^*(2i) \leq (1 - 2\beta + D + \beta D + D^2)(1 - 2\eta + D + 3\eta D + D^2) < \delta_{\infty,11}$$

respectively (in particular this rules out blocks $1, 1$). Hence we can successively rule out $t_{2i}, t_{2i+1} = -3, a$ with $t_{2i+2} = 1$ or with $t_{2i+2} = -1$ else

$$s_2^*(2i) \leq \eta(1 + \beta - 3\beta + aD + \beta D)(1 - \beta - \beta + D + \beta D) < \delta_{\infty,11},$$

$$s_2^*(2i) \leq \eta(1 + \beta - 3\beta + aD)(1 - \beta + \beta + D) < \delta_{\infty,11},$$

and blocks $t_{2i+1}, ... = a, -3, a, -3, 1$ else

$$s_4^*(2i + 3) \leq \eta(1 + \beta - 3\beta + D)(1 - (\beta - 3\beta + aD - 3\beta D)) < \delta_{\infty,11}.$$

Hence either γ has period $F_1 = (a, -1)$ or period $F_3 = (a, -3)$ or the $t_{2i-1} = a$ occur inside blocks $H = (1, -3, a, -3, 1)$, $H' = (-1, 1, a, 1, -1)$ or $\pm 1, -1, a, -1, \pm 1$. Moreover any block H or H' must be followed (and preceded) by a block $G = (-1, a, -1)$ since $t_{2i-1}, ..., t_{2i+3} = a, -3, 1, -1, 1$ leads to $\min\{s_2^*(2i), s_4^*(2i + 1)\} \leq \eta(1 - (\beta - 3\beta + D - \beta D + \frac{D^2 - \beta D^2 + aD^3 - \beta D^3}{1 - D^2})^2)$ and any G must be followed by an H or H' since blocks $t_{2i}, ..., t_{2i+3} = -1, -1, 1, -1$ give

$$s_1^*(2i) \leq (1 - 2\beta + aD - \beta D + D^2)(1 - 2\eta + D - \eta D + D^2) < \delta_{\infty,11}.$$

Hence apart from period B_1 (where $\delta_0 < \delta_{\infty,11}$) or F_2 (or their negatives) we can assume the sequence consists of blocks HG or $H'G$. We suppose that γ is not in S_{-9} so that we have blocks $...|t_{2i-1}... = ...|HGHG...$ and write

$$d_{2i}^- = -3\beta + D - \lambda + D^3 w^-, \quad d_{2i+1}^+ = -3\beta + D + \lambda + D^7 w^+,$$

where $\lambda := \frac{D^3(1-2\beta-2\beta D+D^2)}{1+D^4}$, so that

$$s_2^*(2i) = \eta\left(1-2\beta+D-\lambda+D^3w^-\right)(1+2\beta-D-\lambda-D^7w^+)$$
$$s_4^*(2i+9) = \eta\left(1-2\beta+D-\lambda+D^3w^+\right)(1+2\beta-D-\lambda-D^7w^-).$$

Hence if w^+ and w^- are of opposite signs then one of these is certainly less than $\delta_{\infty,11}$ and if both are negative (say $w^- < w^+$) then $s_2^*(2i) \leq \delta_{\infty,11} + D^3w^-(1-D^4) < \delta_{\infty,11}$. Hence we can assume that $w^-, w^+ > 0$ and hence that we are in a block $...HG(HGH'G)^lHGHG(H'GHG)^kHG....$ Now

$$w^+ \geq 2(1-2\beta-2\beta D+D^2)\frac{D^{8k}(1-D^4-D^8)}{(1-D^8)}, \quad w^- \leq 2(1-2\beta-2\beta D+D^2)\frac{D^{8l}}{(1-D^8)}$$

hence if $l \geq k+1$ we have

$$s_2^*(2i) \leq \delta_{\infty,11} - 2(1-2\beta-2\beta D+D^2)\frac{D^{8k+7}}{1-D^8}\left((1-D^4-D^8)(1-2\beta)-D^4(1+2\beta)\right)$$

less than $\delta_{\infty,11}$, leaving only the elements of $\delta_{k,11}$.

Suppose finally that $(a,b) = (3,6)$. Now if $t_{2i+1} = a$ then (in the same way as before) we can assume that $-2\beta < d_{2i}^-, d_{2i+1}^+ < 0$ and hence that $t_{2i}, ...$ and $t_{2i+2}, ...$ are of the form $0, -1, ...$ or $-2, 1, ...$ or $-2, a, ...$ Now we can rule out $t_{2i} = -4$ else

$$s_1^*(2i) \leq \left(1-5\beta+\frac{D+4\beta D}{1-D}\right)\left(1-\eta+\frac{\eta+4D}{1-D}\right) < \delta_{\infty,12}$$

and blocks $t_{2i+1}, t_{2i} = -1, -2$ else

$$s_1(2i) \leq \left(1-2\eta+\frac{2D+\eta D}{1-D}\right)\left(1-3\beta+\frac{D+2\beta D}{1-D}\right) < \delta_{\infty,12}$$

and $t_{2i+1}, t_{2i+2}, t_{2i+3} = -1, 0, -1$, else

$$s_1^*(2i+1) \leq \left(1-2\eta+\frac{2D-\eta D}{1-D}\right)\left(1-\beta-\frac{(D-2\beta D)}{1-D}\right) < \delta_{\infty,12}.$$

We can also rule out blocks $t_{2i+1}, t_{2i+2}, t_{2i+3} = -1, 0, 1$ else

$$s_1^*(2i+1) = (1-2\eta+\eta d_{2i}^-)(1-\beta+D-D(-d_{2i+2}^+))$$
$$s_3^*(2i+2) = (1-2\eta+\eta(-d_{2i+2}^+))(1-\beta+D-Dd_{2i}^-))$$

with $d_{2i}^- \le (2\beta - D)/(1-D)$ and the smallest of these is at most $(1-2\eta+\eta d_{2i}^-)(1-\beta+D-Dd_{2i}^-) \le \delta_{\infty,12}$. In particular this means that $t_{2i+1}=1$ occur only in blocks $-2,1,-2$. Now if $t_{2i+1},t_{2i+2},t_{2i+3} = a,-2,1$ then $t_{2i},t_{2i-1} = -2,1$ else $d_{2i+1}^+ \le -2\beta + D$ and $d_{2i}^- \ge -2\beta + aD - 2\beta D$ and $s_4^*(2i) \le \eta(1+\beta D)(1-\beta+D) < \delta_{\infty,12}$. Hence either γ or its negative has period $B_2' = (1,-2)$ or $F_2 = (a,-2)$ or takes the form $...|t_{2i+1}... = ...F_2{B_2'}^l|F_2{B_2'}^kF_2...$ with $d_{2i}^-, d_{2i+1}^+ = (-2\beta+D)/(1-D) + w^-, w^+$ where $w^+ \ge (a-1)D^{k+1}$, $w^- \le (a-1)D^{l+1}/(1-D)$ and if $l \ge k+1$ then $s_2^*(2i) \le \delta_{\infty,12} - \eta(a-1)D^{k+1}((1-\beta)-D(1+\beta)/(1-D))$. This leaves only the elements of $S_{k,12}$. ■

6. Proof of Theorem 3 ($a = 2$)

We assume that α has $a_{2i-1} = a = 2$, $a_{2i} = b \ge 5$, and that γ has $M^*(\alpha,\gamma) > \delta_\infty$. If $t_{2i-1} = 0$ then

$$s_1^*(2i-1) = \eta(1-\beta+d_{2i-2}^-)(1-\beta+d_{2i-1}^+)$$
$$s_3^*(2i-1) = \eta(1-\beta-d_{2i-2}^-)(1-\beta-d_{2i-1}^+)$$

and if $d = \max\{\left|d_{2i-2}^-\right|, \left|d_{2i-1}^+\right|\}$ then the minimum of these is certainly at most $\eta((1-\beta)^2 - d^2)$. Hence when b is even we must have $d \le \beta$ ruling out $t_{2i-1\pm1} \ge 2$ or $t_{2i-1\pm1} \le -4$ or $t_{2i-1\pm1} = -2, t_{2i-1\pm2} = 0$ and hence $t_{2i-1\pm1}, t_{2i-1\pm2}, ... = 0, ...$ or $-2, a, ...$ Likewise if b is odd we must have $d \le \frac{1}{2}\beta^2$ ruling out $t_{2i-1\pm1} \ge 1$ or $t_{2i-1\pm1} \le -3$ or $t_{2i-1\pm1}, t_{2i-1\pm2} = -1, 0$ and any $t_{2i-1} = 0$ must occur inside a block $t_{2i-3}, ..., t_{2i+1} = a, -1, 0, -1, a$.

If $t_{2i-1} = a$ then

$$s_2^*(2i-2) = \eta(1+\beta+d_{2i-2}^-)(1-\beta-d_{2i-1}^+)$$
$$s_4^*(2i-1) = \eta(1-\beta-d_{2i-2}^-)(1+\beta+d_{2i-1}^+).$$

We suppose that $t_{2j} \le t$ for all j with $t_{2i} = t$ infinitely often. If $t \ge 1$ then in view of the above we can assume that they occur inside blocks $t_{2i-1}, t_{2i}, t_{2i+1} = a, t, a$ and hence that $t \le b-4$. Since

$$s_2^*(2i-2) = \eta(1-\beta+(2\beta+d_{2i-2}^-))(1-\beta-(t\beta+aD)+2\beta D - D(2\beta+d_{2i+1}^+))$$
$$s_4^*(2i+1) = \eta(1-\beta+(2\beta+d_{2i+1}^+))(1-\beta-(t\beta+aD)+2\beta D - D(2\beta+d_{2i-2}^-)).$$

and if $d = \min\{2\beta + d^-_{2i-2}, 2\beta + d^+_{2i+1}\}$ then certainly

$$\min\{s^*_2(2i-2), s^*_4(2i+1)\} \leq \eta(1-\beta+d)(1-\beta-(t\beta+aD)+2\beta D - Dd).$$

Now if t_{2i-2} or $t_{2i+2} \leq t-2$ then $d \leq (t\beta+aD)/(1-D)$ and, as $(t\beta+aD)/(1-D) \leq 1 \leq (b-t-3+D)/(1+D) = \frac{1}{2}\left((1-\beta-(t\beta+aD)+2\beta D)/D - (1-\beta)\right)$,

$$\begin{aligned}\min\{s^*_2(2i-2), s^*_4(2i+1)\} &\leq \eta\left((1-\beta+\beta D)^2 - \left(\frac{t\beta+aD}{1-D} - \beta D\right)^2\right)\\ &\leq \eta((1-\beta)^2 + 2\beta D - ((t+1)\beta)^2) < \delta_\infty.\end{aligned}$$

Hence either γ has period (a,t) or $t \leq 0$. We assume that γ is not in $S_{0,t}$ (these were dealt with in Lemma 3). Now observing that if γ has blocks $(t_{2i-1}, t_{2i}, t_{2i+1}) = (0,-\lambda,0)$ or $(a,-\lambda,0)$ or $(a,-\lambda,a)$ then $1-\alpha-\gamma$ contains corresponding blocks $(0,\lambda,0)$ or $(a,\lambda-2,0)$ or $(a,\lambda-4,a)$ we readily deduce that our sequence consists of blocks $(a,-1,0,-1,a)$ or $(a,-1,a)$ or $(a,-3,a)$ when b is odd, and when b is even that $t_{2i-1} = a$ are adjoined by 0 or -2 or $(-4,a)$. Now if b is even and $t_{2i-3},...,t_{2i-1} = a,-2,0$ and $t_{2i} = -2$ or $t_{2i}, t_{2i+1} = 0,0$ then

$$\begin{aligned}s^*_2(2i-1) &\leq \eta\left(1-\beta-2\beta+\frac{aD}{1-D}\right)\left(1-\beta+\frac{aD^2}{1-D}\right)\\ &\leq \eta\left((1-\beta)^2 - \beta(1-\beta-3D)\right) < \delta_\infty.\end{aligned}$$

Hence when b is even either γ has period $0,0$ or the $t_{2i-1} = 0$ occur inside blocks $a,0,0,-2,a$ or their negatives $a,-2,0,0,a$. Hence we can assume that the expansion of γ consists entirely of blocks $(a,0,0,-2)$ or $(a,-2,0,0)$ or $(a,-2)$ or $(a,0)$ or $(a,-4)$ when b is even and blocks $(a,-1,0,-1)$ or $(a,-1)$ or $(a,-3)$ when b is odd.

We deal first with b even. Suppose that we have a block $t_{2i-1}, t_{2i}, t_{2i+1} = a,0,a$ then writing $d^-_{2i-2}, d^+_{2i+1} = -\beta + w^-, -\beta + w^+$ (where $-\beta = (-2\beta+aD)/(1-D)$) we have

$$\begin{aligned}s^*_2(2i-2) &= \eta(1+w^-)(1-2\beta-Dw^+)\\ s^*_4(2i+1) &= \eta(1+w^+)(1-2\beta-Dw^-)\end{aligned}$$

Now writing $w = \min\{w^+, w^-\}$ the minimum of these is plainly at most $\eta(1+w)(1-2\beta-Dw)$ and we can assume that $w^+, w^- > 0$. So if $t_{2i-1},...,t_{2i+1} = a,0,a$ then the

adjacent $t_{2i-2}, t_{2i-3}, ...$ or $t_{2i+2}, t_{2i+3}, ...$ take the form $(-2,a)^l\{(0,a)$ or $(0,0,-2,a)\}, ...,$ and if $t_{2i-1}, ..., t_{2i+1} = a, -4, a$, take the form $(-2,a)^l\{(-4,a)$ or $(-2,0,0,a)\}, ...$.

Similarly suppose that we have a block $t_{2i-3}, ..., t_{2i+1} = a, 0, 0, -2, a$ then writing $d^-_{2i-4}, d^+_{2i+1} = -\beta + v^-, -\beta + v^+$ then

$$s_3^*(2i-2) = \eta(1 - Dv^+)(1 - 2\beta - Dv^-)$$
$$s_1^*(2i-1) = \eta(1 + Dv^-)(1 - 2\beta + Dv^+)$$

the previous conditions $|d^-_{2i-2}|, |d^+_{2i-1}| \le \beta$ giving $v^+ \ge 0$ and $v^- \le 0$. Hence this block must be contained inside a block

$$\{(a,-4) \text{ or } (a,0,0,-2)(a,-2)^m(a,0,0,-2,a)(-2,a)^l\{(0,0,-2,a) \text{ or } (0,a)\}$$

Combining these two restrictions it is clear that γ (it or its negative) must consist solely of blocks of the form $(a,0)(a,-2)^{l_i}$ or solely of the form $(a,0,0,-2)(a,-2)^{l_i}$.

Suppose then that we have a block

$$...(a,0)(a,-2)^m(a,0,a)(-2,a)^l,(0,a),...$$

with $m \ge l+1$. Then $w^+ \ge 2\beta D^l$ and $w^- \le 2\beta D^m/(1-D)$ and

$$\begin{aligned} s_2^*(2i-2) &\le \eta\left(1 + \frac{2\beta D^{l+1}}{1-D}\right)(1 - 2\beta - 2\beta D^{l+1}) \\ &\le \eta\left(1 - 2\beta - \frac{2\beta D^{l+1}}{1-D}((1-D) - (1-2\beta))\right) < \delta_\infty. \end{aligned}$$

Likewise if we have blocks

$$...(a,0,0,-2)(a,-2)^m(a,0,0,-2,a)(-2,a)^l(0,0,-2,a)...$$

with $l \ge m+1$ then

$$v^- \le -(aD - 2\beta D)D^m = -\beta(1-D)D^m, \quad v^+ \le (2\beta - aD)D^l/(1-D) \le \beta D^{m+1}$$

and

$$s_1^*(2i-1) \le \eta\left(1 - 2\beta - \beta D^{m+1}((1-D)(1-2\beta) - D)\right) < \delta_\infty.$$

This leaves only the periodic elements of S_{2k} and S_{2k+1}.

Suppose now that b is odd and that we have a block $t_{2i-3},...,t_{2i+1}=a,-1,0,-1,a$. We assume without loss of generality that $d^+_{2i+1}\leq d^-_{2i-4}$ (else reverse their roles) and write

$$d^-_{2i-4}=\frac{-\beta+aD-3\beta D+aD^2}{1-D^2}+w^-,\quad d^+_{2i+1}=\frac{-3\beta+aD-\beta D+aD^2}{1-D^2}+w^+,$$

so that

$$s_1^*(2i-1)=\eta\left(1-\beta-\frac{1}{2}\beta^2+Dw^+\right)\left(1-\beta+\frac{1}{2}\beta^2+Dw^-\right),$$
$$s_3^*(2i-1)=\eta\left(1-\beta-\frac{1}{2}\beta^2-Dw^-\right)\left(1-\beta+\frac{1}{2}\beta^2-Dw^+\right),$$

the previous restriction $|d^-_{2i-2}|,|d^+_{2i-1}|\leq\frac{1}{2}\beta^2$ forcing $-2\beta+\frac{1}{2}\beta^2\leq d^-_{2i-4},d^+_{2i+1}\leq-\frac{1}{2}\beta^2$ and hence $w^-\leq 0$, $w^+\geq 0$. So the sequence must take the form

$$\{(a,-3)\text{ or }(a,-1,0,-1)\}(a,-3,a,-1)^l(a,-1,0,-1,a)(-3,a,-1,a)^m\{(-1,a)\text{ or }(-1,0,-1,a)\}$$

(or its reflection) with

$$w^-\leq\left(-aD+\frac{2\beta D}{1-D^2}\right)D^{2l},\quad w^+\leq\frac{2\beta D^{2m}}{1-D^2}.$$

Hence if $m\geq l+1$ we have

$$s_1^*(2i-1)\leq\delta_\infty+\frac{D^{2l+2}}{1-D^2}\left(2\left(1-\beta+\frac{1}{2}\beta^2\right)D^2-(1-D-D^2-D^3)\left(1-\beta-\frac{1}{2}\beta^2\right)\right)<\delta_\infty.$$

Taking negatives similarly dismisses $l\geq m+1$ so we can assume that $l=m$ for such blocks.

Suppose now that we have a block $t_{2i-3},...,t_{2i+1}=a,-1,a,-1,a$ then writing

$$d^-_{2i-4},d^+_{2i+1}=\frac{-3\beta+aD-\beta D+aD^2}{1-D^2}+v^-,v^+$$

we have

$$s_2^*(2i-4)=\eta\left(1-\beta+\frac{1}{2}\beta^2+v^-\right)\left(1-\beta-\frac{1}{2}\beta^2-D^2v^+\right)$$
$$s_4^*(2i+1)=\eta\left(1-\beta+\frac{1}{2}\beta^2+v^+\right)\left(1-\beta-\frac{1}{2}\beta^2-D^2v^-\right).$$

Now we can assume that $v^-, v^+ > 0$ else $v = \min\{v^+, v^-\} < 0$ and the minimum of these is at most $\eta(1-\beta+\frac{1}{2}\beta^2+v)(1-\beta-\frac{1}{2}\beta^2-D^2v) < \delta_\infty$. In particular we can rule out blocks $(a,-1,0,-1,a)(-3,a,-1,a)^k(-1,a)$ or $(a,-3)(a,-3,a,-1)^k(a,-1,0,-1,a)$ when $k \geq 1$. Hence we can assume that the blocks $(a,-1,0,-1,a)$ only occur in the periodic elements of S_{2k+1}. Since we have also ruled out blocks $(a,-3)$ $(a,-3,a,-1)^k$ $(a,-1)$, $k \geq 1$, it remains only to consider γ in S_{-2} or γ (or its negative) consisting solely of blocks $(-1,a)(-3,a,-1,a)^{l_i}$ with the $l_i \geq 0$. Suppose then that we have blocks

$$....(a,-1)(a,-1,a,-3)^l(a,-1,a,-1,a)(-3,a,-1,a)^m(-1,a)(-3,a,-1,a)^k(-1,a)....$$

Now

$$v^- \geq \left(2\beta - \frac{2\beta D}{(1+D)}\right)D^{2l} = \frac{2\beta D^{2l}(1-D)}{1-D^2},$$

and if $m \geq l+2$

$$v^+ \leq \frac{2\beta D^{2m}}{1-D^2} \leq \frac{2\beta D^{2l+4}}{1-D^2},$$

and if $m = l+1$, $k \geq 1$,

$$v^+ \leq \left(2\beta - 2\beta D + \frac{2\beta D^2}{1-D^2}\right)D^{2m},$$

and in either case

$$v^+ \leq \frac{2\beta D^{2l+2}(1-D+D^3)}{1-D^2},$$

giving

$$s_4^*(2i+1) \leq \delta_\infty + \frac{2D^{2l+3}}{1-D^2}\left((1-D+D^3)\left(1-\beta-\frac{1}{2}\beta^2\right)-(1-D)\left(1-\beta+\frac{1}{2}\beta^2\right)\right) < \delta_\infty.$$

Hence $m \leq l+1$ and if $m = l+1$ we must have $k = 0$ and (since similar reasoning gives $m \leq k+1$) either $l = m$ or $(l,m,k) = (0,1,0)$. But in this latter case looking at the preceding block

$$...(a,-1)(a,-1,a,-3)^r(a,-1)(a,-1,a,-1,a)(-3,a,-1,a)(-1,a)(-1,a),....$$

we must have $r = 0$ or 1. But if $r = 0$ then $v^- \geq 2\beta$, $v^+ \leq 2\beta D^2/(1-D^2)$ and

$$s_4^*(2i+i) \leq \delta_\infty + \frac{2D^3}{1-D^2}\left((1-\beta-\frac{1}{2}\beta^2)-(1-D^2)(1-\beta+\frac{1}{2}\beta^2)\right) < \delta_\infty.$$

Hence this case only occurs when we alternate between blocks $(-1,a)(-3,a,-1,a)^l$ of length $l = 1$ and $l = 0$ giving the elements of S_{-1}. The remaining γ have blocks of equal length giving the elements of S_{2k}. ■

References

[**1**] E. S. Barnes & H. P. F. Swinnerton-Dyer, *The inhomogeneous minima of binary quadratic forms.* Part I, *Acta Math.* **87** (1952), 259-323, Part II, *Acta M ath.* **88** (1952), 279-316, Part III, *Acta Math.* **92** (1 954), 199-234, Part IV (without second author) *Acta Math.* **92** (1954), 235-2 64..

[**2**] D. Cardon, *A Euclidean ring containing* $\mathbb{Z}[\sqrt{14}]$, C. R. Math. Rep. Acad. Sci. Canada **19** (1997), 28–32.

[**3**] D. J. Crisp, W. Moran & A. D. Pollington, *Inhomogeneous diophantine approximation and Hall's ray preprint.*

[**4**] T. W. Cusick and M. E. Flahive, *The Markoff and Lagrange spectra, mathematical surveys and monographs, no. 30, American Mathematical Society*, 1989.

[**5**] H. J. Godwin, *On a conjecture of Barnes and Swinnerton-Dyer*, Proc. Camb. Phil. Soc. **59** (1963), 519–522.

[**2**] M. Harper, *Reference??.*

[**5**] F. Lemmermeyer, *The Euclidean algorithm in algebraic number fields*, Expo. Math. **13** (1995), 385–416.

[**6**] C. G. Pinner, *More on inhomogeneous Diophantine approximation*, preprint.

[**7**] P. Varnavides, *Non-homogeneous quadratic forms, I*, Proc. K. Ned. Akad. Wet. (Indag. Math.) **51** (1948), 396–404, 470–481.

Mathematics and Computer Science, University of Northern British Columbia, 3333 University Way, Prince George, B.C., Canada V2N 4Z9. pinner@math.ksu.edu